

## Datasheet: STAR117F

**BATCH NUMBER 168696**

<b>Description:</b>	GOAT ANTI MOUSE IgG (H/L):FITC (MULTI SPECIES ADSORBED)
<b>Specificity:</b>	IgG (H/L)
<b>Format:</b>	FITC
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.5 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/50 - 1/200
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
Immunofluorescence			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

<b>Target Species</b>	Mouse						
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
<b>Max Ex/Em</b>	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					
<b>Preparation</b>	Purified IgG prepared by affinity chromatography.						
<b>Antiserum Preparation</b>	Antisera to mouse IgG were raised by repeated immunizations of goats with highly purified antigen.						
<b>Buffer Solution</b>	Phosphate buffered saline.						
<b>Preservative</b>	0.09% Sodium Azide						
<b>Stabilisers</b>	0.2% Bovine Serum Albumin						

<b>Approx. Protein Concentrations</b>	IgG concentration 0.5 mg/ml.
<b>Immunogen</b>	Whole mouse IgG
<b>External Database Links</b>	<p><b>UniProt:</b></p> <p><a href="#">P01837</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01869</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01867</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01864</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01843</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01865</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01844</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01868</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01724</a>    <a href="#">Related reagents</a></p> <p><a href="#">P03987</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01863</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01845</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">16071</a>    Igk-C        <a href="#">Related reagents</a></p> <p><a href="#">16017</a>    Ighg1        <a href="#">Related reagents</a></p> <p><a href="#">16016</a>    Ighg2b      <a href="#">Related reagents</a></p> <p><a href="#">380793</a>    Igh-1a      <a href="#">Related reagents</a></p> <p><a href="#">380793</a>    Igh-1a      <a href="#">Related reagents</a></p> <p><a href="#">433053</a>    LOC433053   <a href="#">Related reagents</a></p> <p><a href="#">16017</a>    Ighg1        <a href="#">Related reagents</a></p> <p><a href="#">16142</a>    Iglv1        <a href="#">Related reagents</a></p> <p><a href="#">110786</a>    Iglc2        <a href="#">Related reagents</a></p> <p><a href="#">110787</a>    Iglc3        <a href="#">Related reagents</a></p> <p><a href="#">380793</a>    Igh-1a      <a href="#">Related reagents</a></p> <p><a href="#">380795</a>    AI324046    <a href="#">Related reagents</a></p>
<b>Synonyms</b>	Igh-4
<b>RRID</b>	AB_324190
<b>Specificity</b>	<p><b>Goat anti Mouse IgG antibody</b> recognizes mouse IgG and light chains common to other mouse immunoglobulin classes.</p> <p>Goat anti Mouse IgG has been cross-adsorbed using human, bovine, porcine, equine, lapine and chicken immunoabsorbants to remove cross-reactive antibodies. Less than 0.1% cross reactivity was detected to human, bovine, porcine, equine, caprine, lapine and chicken IgG by immunoelectrophoresis and ELISA.</p>

Goat anti Mouse IgG antibody is highly recommended for use as a secondary antibody with human and veterinary samples. Goat anti Mouse IgG antibody has been used successfully as a secondary detection reagent in combination with mouse clone [CC327](#) for the detection of TNF $\alpha$  and mouse clone [8M6](#) for the detection of interleukin-8 in bovine respiratory syncytial virus infected, neonatal ovine lung tissue by immunohistochemistry ([Redondo et al. 2013](#)).

---

**Flow Cytometry** Use 50  $\mu$ l of the suggested working dilution to label  $1 \times 10^6$  cells in 100  $\mu$ l

---

- References**
1. Banerjee, K. *et al.* (2012) Occluding the mannose moieties on human immunodeficiency virus type 1 gp120 with griffithsin improves the antibody responses to both proteins in mice. [AIDS Res Hum Retroviruses. 28 \(2\): 206-14.](#)
  2. Abdala-Valencia, H. *et al.* (2012) Vitamin E isoforms differentially regulate intercellular adhesion molecule-1 activation of PKC $\alpha$  in human microvascular endothelial cells. [PLoS One. 7: e41054.](#)
  3. Redondo, E. *et al.* (2014) Induction of interleukin-8 and interleukin-12 in neonatal ovine lung following experimental inoculation of bovine respiratory syncytial virus. [J Comp Pathol. 150 \(4\): 434-48.](#)
  4. Askari, N. *et al.* (2015) Tetracycline-regulated expression of OLIG2 gene in human dental pulp stem cells lead to mouse sciatic nerve regeneration upon transplantation. [Neuroscience. 305: 197-208.](#)
  5. Iwaszko-Simonik, A. *et al.* (2015) Expression of surface platelet receptors (CD62P and CD41/61) in horses with recurrent airway obstruction (RAO). [Vet Immunol Immunopathol. 164 \(1-2\): 87-92.](#)
  6. Singh, S.M. *et al.* (2016) Characterization of Immune Responses to an Inactivated Avian Influenza Virus Vaccine Adjuvanted with Nanoparticles Containing CpG ODN. [Viral Immunol. 29 \(5\): 269-75.](#)
  7. Alimolaei, M. *et al.* (2017) A Recombinant Probiotic, *Lactobacillus casei*, Expressing the *Clostridium perfringens*  $\alpha$ -toxoid, as an Orally Vaccine Candidate Against Gas Gangrene and Necrotic Enteritis. [Probiotics Antimicrob Proteins. Apr 11 \[Epub ahead of print\].](#)
  8. Topoluk, N. *et al.* (2017) Amniotic Mesenchymal Stromal Cells Exhibit Preferential Osteogenic and Chondrogenic Differentiation and Enhanced Matrix Production Compared With Adipose Mesenchymal Stromal Cells. [Am J Sports Med. 45 \(11\): 2637-46.](#)
  9. Schmidli, M.R. *et al.* (2018) Inflammatory pattern of the infrapatellar fat pad in dogs with canine cruciate ligament disease. [BMC Vet Res. 14 \(1\): 161.](#)
  10. Li, T. *et al.* (2021) RNF167 activates mTORC1 and promotes tumorigenesis by targeting CASTOR1 for ubiquitination and degradation. [Nat Commun. 12 \(1\): 1055.](#)
  11. Dicks, M.D.J. *et al.* (2022) Modular capsid decoration boosts adenovirus vaccine-induced humoral immunity against SARS-CoV-2. [Mol Ther. 30 \(12\): 3639-57.](#)
  12. Soleimani, M. *et al.* (2022) Covalent JNK Inhibitor, JNK-IN-8, Suppresses Tumor Growth in Triple-Negative Breast Cancer by Activating TFEB- and TFE3-Mediated Lysosome Biogenesis and Autophagy. [Mol Cancer Ther. 21 \(10\): 1547-60.](#)

---

**Storage** Store at +4°C. DO NOT FREEZE.  
This product should be stored undiluted. This product is photosensitive and should be protected from light.  
Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch.

---

**Health And Safety Information** Material Safety Datasheet documentation #10041 available at:  
<https://www.bio-rad-antibodies.com/SDS/STAR117F>  
10041

---

**Regulatory** For research purposes only.

---

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M428646:240301'

**Printed on 26 Jun 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)