

Datasheet: STAR121F BATCH NUMBER 153257

Description:	GOAT ANTI RABBIT IgG (Fc):FITC
Specificity:	IgG (Fc)
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/200 - 1/2000
Immunohistology - Frozen				
Immunohistology - Paraffin			•	
ELISA			•	
Immunoprecipitation			•	
Western Blotting			•	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

	1110	430	320	
	FITC	490	525	
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid			
Target Species	Rabbit			

Antiserum Preparation Antisera to rabbit IgG were raised by repeated immunisations of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
	0.2% Bovine Serum Albumin

IgG concentration 1.0 mg/ml	
UniProt: P01870 Related reagents	
Entrez Gene: 100009097 LOC100009097 Related reagents	
AB_567028	
Goat anti rabbit IgG antibody recognizes the Fc region of rabbit im confirmed by ELISA and immunoelectrophoresis. The antibody has recross adsorbed and may cross react with IgG from other species. It recognition of rabbit IgM in ELISA.	not been species
Goat anti rabbit IgG antibody has been used successfully as a secon reagent for rabbit primary antibodies using immunohistochemical ted <u>al. 2013</u>).	•
 Kalluri, S.R. <i>et al.</i> (2010) Quantification and functional characterize native aquaporin 4 in neuromyelitis optica. <u>Arch Neurol. 67:1201-8.</u> Redondo, E. <i>et al.</i> (2014) Induction of interleukin-8 and interleukin lung following experimental inoculation of bovine respiratory syncytiate Pathol. 150 (4): 434-48. 	n-12 in neonatal ovine
3. Tyrsina, E. <i>et al.</i> (2019) Detection and quantification of VEGFR-1 cells by a new flow cytometry-based method. <u>J Immunotoxicol. 16 (1</u> 4. Lin, W. <i>et al.</i> (2020) Rapid identification of anti-idiotypic mAbs with diverse epitopes by rabbit single B-cell sorting-culture and cloning te <u>15 (12): e0244158.</u>	l <u>): 74-81.</u> n high affinity and
5. Li, Y. <i>et al.</i> (2022) Low-Temperature Plasma-Activated Medium Infand Progression of Lung Cancer by Targeting the PI3K/Akt and MAF Med Cell Longev. 2022: 9014501.	
Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensiti protected from light. Should this product contain a precipitate we rec microcentrifugation before use.	
12 months from date of despatch	
Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/STAR121F 10041	
For research purposes only	
	Entrez Gene: 100009097 LOC100009097 Related reagents AB_567028 Goat anti rabbit IgG antibody recognizes the Fc region of rabbit im confirmed by ELISA and immunoelectrophoresis. The antibody has a cross adsorbed and may cross react with IgG from other species. It recognition of rabbit IgM in ELISA. Goat anti rabbit IgG antibody has been used successfully as a secon reagent for rabbit primary antibodies using immunohistochemical ted al. 2013). 1. Kalluri, S.R. et al. (2010) Quantification and functional characteriz native aquaporin 4 in neuromyelitis optica. Arch Neurol. 67:1201-8. 2. Redondo, E. et al. (2014) Induction of interleukin-8 and interleukin lung following experimental inoculation of bovine respiratory syncytic Pathol. 150 (4): 434-48. 3. Tyrsina, E. et al. (2019) Detection and quantification of VEGFR-1 cells by a new flow cytometry-based method. J Immunotoxicol. 16 (4). Lin, W. et al. (2020) Rapid identification of anti-idiotypic mAbs with diverse epitopes by rabbit single B-cell sorting-culture and cloning to 15 (12): e0244158. 5. Li, Y. et al. (2022) Low-Temperature Plasma-Activated Medium In and Progression of Lung Cancer by Targeting the PI3K/Akt and MARMed Cell Longev. 2022: 9014501. Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensit protected from light. Should this product contain a precipitate we recomic product should be stored undiluted. This product is photosensit protected from light. Should this product contain a precipitate we recomic product should be stored undiluted. This product is photosensit protected from light. Should this product contain a precipitate we recomic product should be stored undiluted. This product is photosensit protected from light. Should this product contain a precipitate we recomic product should be stored undiluted. This product is photosensit protected from light. Should this product contain a precipitate we recomic product should be stored undiluted. This product is photosensit product.

North & South Tel: +1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe Tel: +49 (0) 89 8090 95 21 America

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.comd a Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com

То

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M369650:200529'

Printed on 10 Feb 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint