

Datasheet: STAR121P

| | |
|----------------------|-------------------------------|
| Description: | GOAT ANTI RABBIT IgG (Fc):HRP |
| Specificity: | IgG (Fc) |
| Format: | HRP |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|----------------------|
| Immunohistology - Frozen | | | ▪ | |
| Immunohistology - Paraffin | | | ▪ | |
| ELISA | ▪ | | | 1/10,000 - 1/100,000 |
| Western Blotting | ▪ | | | 1/ 5,000 - 1/ 50,000 |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------------|--|
| Target Species | Rabbit |
| Product Form | Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid |
| Antiserum Preparation | Antisera to rabbit IgG were raised by repeated immunisations of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. |
| Buffer Solution | Phosphate buffered saline |
| Preservative | 0.05% Proclin™ 300 |
| Stabilisers | 0.2% Bovine Serum Albumin |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml |
| External Database Links | <p>UniProt: P01870 Related reagents</p> <p>Entrez Gene: 100009097 LOC100009097 Related reagents</p> |
| RRID | AB_567025 |

| | |
|--------------------------------------|---|
| Specificity | <p>Goat anti rabbit IgG antibody recognizes the Fc region of rabbit immunoglobulin G, confirmed by ELISA and immunoelectrophoresis. The antibody has not been species cross adsorbed and may cross react with IgG from other species. It shows weak recognition of rabbit IgM in ELISA.</p> <p>Goat anti rabbit IgG antibody has been used successfully as a secondary detection reagent for rabbit primary antibodies using immunohistochemical techniques (Redondo et al. 2013).</p> |
| References | <ol style="list-style-type: none"> 1. Kalluri, S.R. <i>et al.</i> (2010) Quantification and functional characterization of antibodies to native aquaporin 4 in neuromyelitis optica. Arch Neurol. 67:1201-8. 2. Redondo, E. <i>et al.</i> (2014) Induction of interleukin-8 and interleukin-12 in neonatal ovine lung following experimental inoculation of bovine respiratory syncytial virus. J Comp Pathol. 150 (4): 434-48. |
| Storage | <p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.</p> |
| Guarantee | 12 months from date of despatch |
| Acknowledgements | Proclin™ 300 is a trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow. |
| Health And Safety Information | <p>Material Safety Datasheet documentation #20391 available at: 20391: https://www.bio-rad-antibodies.com/uploads/MSDS/20391.pdf</p> |
| Regulatory | For research purposes only |

Related Products

Recommended Useful Reagents

[TMB CORE \(BUF056A\)](#)

[TMB CORE+ \(BUF062A\)](#)

[TMB SIGNAL+ \(BUF054A\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M369651:200529'

Printed on 11 Aug 2020