

## Datasheet: STAR122F

**BATCH NUMBER 173116**

<b>Description:</b>	RABBIT ANTI GOAT IgG (Fc):FITC
<b>Specificity:</b>	IgG (Fc)
<b>Format:</b>	FITC
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/200 - 1/2,000
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA			▪	
Immunoprecipitation			▪	
Western Blotting			▪	

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Goat		
<b>Product Form</b>	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
<b>Max Ex/Em</b>	<b>Fluorophore</b>	<b>Excitation Max (nm)</b>	<b>Emission Max (nm)</b>
	FITC	490	525

**Antiserum Preparation** Antisera to goat IgG were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

**Buffer Solution** Phosphate buffered saline

**Preservative Stabilisers** 0.09% Sodium Azide (NaN<sub>3</sub>)

<b>Approx. Protein Concentrations</b>	IgG concentration 1.0 mg/ml
<b>RRID</b>	AB_567029
<b>Specificity</b>	<p><b>Rabbit anti goat IgG</b> recognizes an epitope localised to the Fc region of goat immunoglobulin G as assessed by immunoelectrophoresis and ELISA.</p> <p>Rabbit anti goat IgG may cross react with IgG from other species.</p>
<b>References</b>	<p>1. Corrales, L. <i>et al.</i> (2012) Anaphylatoxin C5a creates a favorable microenvironment for lung cancer progression. <a href="#">J Immunol. 189 (9): 4674-83.</a></p> <p>2. Lam, D.C. <i>et al.</i> (2015) S-maltoheptaose targets syndecan-bound effectors to reduce smoking-related neutrophilic inflammation. <a href="#">Sci Rep. 5: 12945.</a></p>
<b>Storage</b>	<p>This product is shipped at ambient temperature.</p> <p>Store at +4°C. DO NOT FREEZE.</p> <p>This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	12 months from date of despatch
<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10040 available at: <a href="https://www.bio-rad-antibodies.com/SDS/STAR122F">https://www.bio-rad-antibodies.com/SDS/STAR122F</a>
<b>Regulatory</b>	For research purposes only

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)  
'M441618:250523'

**Printed on 19 Mar 2026**