

## Datasheet: STAR122P

**BATCH NUMBER 172758**

<b>Description:</b>	RABBIT ANTI GOAT IgG (Fc):HRP
<b>Specificity:</b>	IgG (Fc)
<b>Format:</b>	HRP
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen	▪			1/200 - 1/5000
Immunohistology - Paraffin			▪	
ELISA	▪			1/ 10,000 - 1/ 100,000
Immunoprecipitation			▪	
Western Blotting	▪			1/5000 - 1/50000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

**Target Species** Goat

**Product Form** Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid

**Antiserum Preparation** Antisera to goat IgG were raised by repeated immunisations of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

**Buffer Solution** Phosphate buffered saline.

**Preservative Stabilisers** 0.05% ProClin 300

**Approx. Protein Concentrations** IgG concentration 1.0 mg/ml.

RRID AB\_567020

---

**Specificity** Rabbit anti goat IgG recognizes an epitope localised to the Fc region of goat immunoglobulin G as assessed by immunoelectrophoresis and ELISA.

Rabbit anti goat IgG may cross react with IgG from other species.

---

**References**

1. Corrales, L. *et al.* (2012) Anaphylatoxin C5a creates a favorable microenvironment for lung cancer progression. [J Immunol. 189 \(9\): 4674-83.](#)
2. Lam, D.C. *et al.* (2015) S-maltoheptaose targets syndecan-bound effectors to reduce smoking-related neutrophilic inflammation. [Sci Rep. 5: 12945.](#)

---

**Storage** This product is shipped at ambient temperature.  
Store at +4°C. DO NOT FREEZE.  
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.

---

**Guarantee** 12 months from date of despatch.

---

**Acknowledgements** ProClin is a trademark of The Dow Chemical Company (“Dow”) or an affiliated company of Dow.

---

**Health And Safety Information** Material Safety Datasheet documentation #20392 available at: <https://www.bio-rad-antibodies.com/SDS/STAR122P>

---

**Regulatory** For research purposes only.

---

## Related Products

### Recommended Useful Reagents

[AbGUARD® HRP STABILIZER PLUS \(BUF052A\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052B\)](#)

[AbGUARD® HRP STABILIZER PLUS \(BUF052C\)](#)

[TMB CORE \(BUF056A\)](#)

[TMB CORE+ \(BUF062A\)](#)

[TMB SIGNAL+ \(BUF054A\)](#)

**Product inquiries:** [www.bio-rad-antibodies.com/technical-support](http://www.bio-rad-antibodies.com/technical-support)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)

'M441619:250523'

Printed on 18 Dec 2025

---