

Datasheet: STAR132F

Description:	GOAT ANTI MOUSE IgG1:FITC
Specificity:	IgG1
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/1000
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
Immunofluorescence	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse						
Species Cross Reactivity	Does not react with:Human						
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid						
Max Ex/Em	<table border="1"> <thead> <tr> <th>Fluorophore</th> <th>Excitation Max (nm)</th> <th>Emission Max (nm)</th> </tr> </thead> <tbody> <tr> <td>FITC</td> <td>490</td> <td>525</td> </tr> </tbody> </table>	Fluorophore	Excitation Max (nm)	Emission Max (nm)	FITC	490	525
Fluorophore	Excitation Max (nm)	Emission Max (nm)					
FITC	490	525					

Antiserum Preparation Antisera to mouse IgG1 were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0mg/ml
Immunogen	Mouse IgG1 paraproteins.

**External Database
Links**

UniProt:

[P01869](#) [Related reagents](#)
[P01868](#) [Related reagents](#)

Entrez Gene:

[16017](#) Ighg1 [Related reagents](#)
[16017](#) Ighg1 [Related reagents](#)

Synonyms

Igh-4

RRID

AB_2124271

Specificity

Goat anti Mouse IgG1 antibody recognizes Mouse IgG1. This antibody has been cross absorbed against mouse IgM, IgG2a, IgG2b, IgG3 and IgA, pooled human sera and purified human paraproteins. Goat anti Mouse IgG1 antibody shows minimal cross-reactivity with human immunoglobulins.

Flow Cytometry

Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Croft, N.P. *et al.* (2009) Stage-specific inhibition of MHC class I presentation by the Epstein-Barr virus BNLF2a protein during virus lytic cycle. [PLoS Pathog. 5\(6\): e1000490.](#)
2. Zuo, J. *et al.* (2011) The Epstein-Barr virus-encoded BILF1 protein modulates immune recognition of endogenously processed antigen by targeting MHC class I molecules trafficking on both the exocytic and endocytic pathways. [J Virol. 85: 1604-14.](#)
3. Knipping, K. *et al.* (2011) A gastrointestinal rotavirus infection mouse model for immune modulation studies. [Virology 438: 109.](#)
4. Young, D. *et al.* (2012) Soy-derived di- and tripeptides alleviate colon and ileum inflammation in pigs with dextran sodium sulfate-induced colitis. [J Nutr. 142 \(2\): 363-8.](#)
5. Bagai, U. and Pawar, A. (2013) A blood stage fraction of *Plasmodium berghei* induces protective and long lasting immune response in BALB/c mice. [Parasitol Int. 62: 329-36.](#)
6. Anda, S. *et al.* (2014) Cell-cycle analyses using thymidine analogues in fission yeast. [PLoS One. 9 \(2\): e88629.](#)
7. Kamat, M.M. *et al.* (2016) Changes in Myeloid Lineage Cells in the Uterus and Peripheral Blood of Dairy Heifers During Early Pregnancy. [Biol Reprod. Aug 10. pii: biolreprod.116.141069. \[Epub ahead of print\]](#)
8. Ramanathan, R. *et al.* (2015) Transplantation of human stem cell-derived hepatocytes in an animal model of acute liver failure. [Surgery. 158 \(2\): 349-59.](#)
9. Hwang, S.R. *et al.* (2015) Altered expression levels of neurodevelopmental proteins in fetal brains of BTBR T+tf/J mice with autism-like behavioral characteristics. [J Toxicol Environ Health A. 78 \(8\): 516-23.](#)
10. Zhao, Z. *et al.* (2015) Multiple B-cell epitope vaccine induces a Staphylococcus enterotoxin B-specific IgG1 protective response against MRSA infection. [Sci Rep. 5: 12371.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product

contain a precipitate we recommend microcentrifugation before use.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M369676:200529'

Printed on 07 Sep 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)