

Datasheet: STAR135F BATCH NUMBER 163224

Description:	GOAT ANTI MOUSE IgG2c:FITC
Specificity:	lgG2c
Other names:	lgh-1b ALLELE
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Immunohistology - Frozen				
Immunohistology - Paraffin			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using the appropriate negative/positive controls.

Target Species	Mouse	Mouse				
Product Form	Purified IgG conju	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer				
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm	n)		
	FITC	490	525			
Preparation	Purified IgG prepa	Purified IgG prepared by affinity chromatography				
Antiserum Preparat	t ion Antisera to mouse purified antigen.	e IgG2c were raised by re	peated immunisatio	ns of goats with highly		
Buffer Solution	Phosphate buffere	ed saline				
Preservative Stabilisers	<0.1% Sodium Az	zide (NaN ₃)				

Approx. Protein Concentrations	IgG concentration 1.0mg/ml				
Immunogen	Native mouse IgG2c.				
RRID	AB_1102667				
Specificity	Goat anti Mouse IgG2c antibody recognizes mouse IgG2c, otherwise known as the Igh 1b allele of IgG2a. Goat anti Mouse IgG2c antibody is assessed by ELISA and immunoelectrophoresis and has been shown to react with IgG2c in C57BL/6, SJL, C57BL/10, CB20, C57BL/6 by Balb/C crosses and pools of serum of outbred mice and does not react with L-chains or other IgG subclasses.				
Flow Cytometry	Use 10 ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.				
References	 Jalili, R.B. et al. (2010) Local Expression of Indoleamine 2,3 Dioxygenase in Syngeneic Fibroblasts Significantly Prolongs Survival of an Engineered Three-Dimensional Islet Allograft. Diabetes 59: 2219-27 Arnold, I.C. et al. (2011) The C-terminally encoded, MHC class II-restricted T cell antigenicity of the Helicobacter pylori virulence factor CagA promotes gastric preneoplasia. J Immunol. 186: 6165-72. Paes, W. et al. (2016) Recombinant polymorphic membrane protein D in combination with a novel, second-generation lipid adjuvant protects against intra-vaginal Chlamydia trachomatis infection in mice. Vaccine. 34 (35): 4123-31. Carroll, E.C. et al. (2016) The Vaccine Adjuvant Chitosan Promotes Cellular Immunity via DNA Sensor cGAS-STING-Dependent Induction of Type I Interferons. Immunity. 44 (3): 597-608. Zouggari, Y. et al. (2013) B lymphocytes trigger monocyte mobilization and impair heart function after acute myocardial infarction. Nat Med. 19 (10): 1273-80. Müller-Winkler, J. et al. (2021) Critical requirement for BCR, BAFF, and BAFFR in memory B cell survival. J Exp Med. 218 (2) [Epub ahead of print]. McEntee, C.P. et al. (2020) Type I IFN signalling is required for cationic adjuvant 				
	formulation (CAF)01-induced cellular immunity and mucosal priming. <u>Vaccine. 38 (3):</u> 635-43.				
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.				
Guarantee	12 months from date of despatch				
Health And Safety Information	Material Safety Datasheet documentation #10041 available at: https://www.bio-rad-antibodies.com/SDS/STAR135F 10041				
Regulatory	For research purposes only				

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

Fax: +1 919 878 3751 Fax: +44 (0)1865 852 739 Fax: +49 (0) 89 8090 95 50 To Email: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio-rad.com Email: antibody_sales_de@bio-rad.com Email: antibody_sales_de@bio-rad.com

batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M389792:210806'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint