

Datasheet: STAR137 **BATCH NUMBER 155307**

Description:	GOAT ANTI MOUSE IgA
Specificity:	IgA
Format:	Purified
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.biorad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			•	
Immunohistology - Frozen			•	
Immunohistology - Paraffin				
ELISA	•			
Immunoprecipitation			•	
Western Blotting			•	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse
Product Form	Purified IgG - liquid

Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Borate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0mg/ml

Immunogen	Mouse IgA paraproteins.
External Database Links	UniProt: P01878 Related reagents Entrez Gene: 16061 Igh-VJ558 Related reagents
RRID	AB_2075638
Specificity	Goat anti Mouse IgA antibody recognizes Mouse IgA and has been cross absorbed against mouse IgM, IgG1, IgG2a, IgG2b and IgG3.
References	 Capoccia, B.J. <i>et al.</i> (2011) Transcription factor MIST1 in terminal differentiation of mouse and human plasma cells. Physiol Genomics. 43 (3): 174-86. Yamashita, H. <i>et al</i> (2012) Overcoming food allergy through acquired tolerance conferred by transfer of Tregs in a murine model. Allergy. 67: 201-9. Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. J Nat Med. 71 (1): 68-75. Ramstead, A.G. <i>et al.</i> (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary <i>Coxiella burnetii</i> Infection. Infect Immun. 84 (4): 940-9.
Storage	Store at +4°C or at -20°C if preferred. This product should be stored undiluted.
	Storage in frost free freezers is not recommended. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.
Guarantee	12 months from date of despatch
Health And Safety Information	Material Safety Datasheet documentation #10077 available at: https://www.bio-rad-antibodies.com/SDS/STAR137 10077
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M369689:200529'

Printed on 30 Jan 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint