

Datasheet: STAR137

Description:	GOAT ANTI MOUSE IgA
Specificity:	IgA
Format:	Purified
Product Type:	Polyclonal Antibody
lsotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio- rad-antibodies.com/protocols. Flow Cytometry • Immunohistology - Frozen • Immunohistology - Paraffin • ELISA • Immunohistology - Paraffin • Western Blotting • Western Blotting • Western Blotting • Western Blotting • Product Form Purified IgG - liquid Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Proservative <0.1% Sodium Azide (NaN ₃) Approx. Protein Concentration 1.0mg/ml	Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal									
Information. For general protocol recommendations, please visit www.bio- rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution Flow Cytometry - - - Immunohistology - Frozen - - - Immunohistology - Paraffin - - - ELISA - - - - Immunoprecipitation - - - - Western Blotting - - - - - Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. Target Species Mouse - - Product Form Purified IgG - liquid - - Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Borate buffered saline - Preservative <0.1% Sodium Azide (NaN ₃) - Approx. Protein IgG concentration 1.0mg/ml - <th></th> <td colspan="8"></td>											
rad-antibodies.com/protocols. Yes No Not Determined Suggested Dilution Flow Cytometry • • • Immunohistology - Prozen • • • Immunohistology - Paraffin • • • Immunohistology - Paraffin • • • Immunoprecipitation • • • Western Blotting • • • Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. Target Species Mouse Product Form Purified IgG - liquid Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Borate buffered saline • Preservative <0.1% Sodium Azide (NaN ₃) • Approx. Protein IgG concentration 1.0mg/ml •		-									
YesNoNot DeterminedSuggested DilutionFlow Cytometry••Immunohistology - Frozen•Immunohistology - Parafini•ELISA•ELISA•Immunoprecipitation•Western Blotting•Western Blotting•Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures. It is recommended that the user titrates the antibody for use in such procedures.Frage SpeciesMouseProduct FormPurified IgG - liquidBuffer SolutionBorate buffered salineFreservative<											
Immunohistology - Frozen • Immunohistology - Paraffin • ELISA • Immunoprecipitation • Western Blotting • Western Blotting • Western Blotting • Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. Target Species Mouse Product Form Purified IgG - liquid Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Borate buffered saline Preservative stabilisers <0.1% Sodium Azide (NaN ₃) Approx. Protein IgG concentration 1.0mg/ml											
Immunohistology - Paraffin • ELISA • Immunoprecipitation • Western Blotting • Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. Target Species Mouse Product Form Purified IgG - liquid Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Borate buffered saline Preservative Stabilisers <0.1% Sodium Azide (NaN ₃) Approx. Protein IgG concentration 1.0mg/ml		Flow Cytometry			•						
ELISA • Immunoprecipitation • Western Blotting • Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls. Target Species Mouse Product Form Purified IgG - liquid Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Borate buffered saline Preservative Stabilisers <0.1% Sodium Azide (NaN ₃) Approx. Protein IgG concentration 1.0mg/ml		Immunohistology - Frozen			•						
Immunoprecipitation•Western Blotting•Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.Target SpeciesMouseProduct FormPurified IgG - liquidAntiserum Preparation antigen. Purified IgG was prepared from whole serum by affinity chromatography.Buffer SolutionBorate buffered salinePreservative Stabilisers<0.1% Sodium Azide (NaN3)Approx. Protein ConcentrationsIgG concentration 1.0mg/ml		Immunohistology - Paraffin			•						
Western BlottingWhere this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.Target SpeciesMouseProduct FormPurified IgG - liquidAntiserum Preparation antigen. Purified IgG was prepared from whole serum by affinity chromatography.Buffer SolutionBorate buffered salinePreservative Stabilisers<0.1% Sodium Azide (NaN3)Approx. Protein ConcentrationsIgG concentration 1.0mg/ml		ELISA	•								
Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.Target SpeciesMouseProduct FormPurified IgG - liquidAntiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.Buffer SolutionBorate buffered salinePreservative Stabilisers<0.1% Sodium Azide (NaN3)		Immunoprecipitation									
necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.Target SpeciesMouseProduct FormPurified IgG - liquidAntiserum PreparationAntisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.Buffer SolutionBorate buffered salinePreservative Stabilisers<0.1% Sodium Azide (NaN3)		Western Blotting			-						
the antibody for use in their own system using appropriate negative/positive controls.Target SpeciesMouseProduct FormPurified IgG - liquidAntiserum PreparationAntisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.Buffer SolutionBorate buffered salinePreservative Stabilisers<0.1% Sodium Azide (NaN_3)		Where this antibody has	not been	tested for	use in a particular tec	hnique this does not					
Target SpeciesMouseProduct FormPurified IgG - liquidAntiserum PreparationAntisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.Buffer SolutionBorate buffered salinePreservative Stabilisers<0.1% Sodium Azide (NaN3)		necessarily exclude its us	se in such	procedur	es. It is recommended	that the user titrates					
Product FormPurified IgG - liquidAntiserum PreparationAntisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.Buffer SolutionBorate buffered salinePreservative Stabilisers<0.1% Sodium Azide (NaN3)		the antibody for use in the	eir own sy	stem usin	g appropriate negativ	e/positive controls.					
Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Borate buffered saline Preservative Stabilisers <0.1% Sodium Azide (NaN ₃) Approx. Protein Concentrations IgG concentration 1.0mg/ml	Target Species	Mouse									
antigen. Purified IgG was prepared from whole serum by affinity chromatography. Buffer Solution Borate buffered saline Preservative Stabilisers <0.1% Sodium Azide (NaN ₃) Approx. Protein Concentrations IgG concentration 1.0mg/ml	Product Form	Purified IgG - liquid									
Preservative Stabilisers <0.1% Sodium Azide (NaN ₃) Approx. Protein Concentrations IgG concentration 1.0mg/ml	Antiserum Preparatio	-			-	•					
Stabilisers <0.1% Sodium Azide (NaN ₃) Approx. Protein Concentrations IgG concentration 1.0mg/ml	Buffer Solution	Borate buffered saline									
Concentrations		<0.1% Sodium Azide (NaN ₃)									
		IgG concentration 1.0mg/ml									
Immunogen Mouse IgA paraproteins.	Immunogen	Mouse IgA paraproteins.									

External Database Links RRID Specificity	UniProt: P01878 Related reagents Entrez Gene: 16061 Igh-VJ558 Related reagents AB_2075638 Goat anti Mouse IgA antibody recognizes Mouse IgA and has been cross absorbed against mouse IgM, IgG1, IgG2a, IgG2b and IgG3.			
References	 Capoccia, B.J. <i>et al.</i> (2011) Transcription factor MIST1 in terminal differentiation of mouse and human plasma cells. <u>Physiol Genomics. 43 (3): 174-86.</u> Yamashita, H. <i>et al</i> (2012) Overcoming food allergy through acquired tolerance conferred by transfer of Tregs in a murine model. <u>Allergy. 67: 201-9.</u> Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. <u>J Nat Med. 71 (1): 68-75.</u> Ramstead, A.G. <i>et al.</i> (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary <i>Coxiella burnetii</i> Infection. <u>Infect Immun. 84 (4): 940-9.</u> Sessevmez, M. <i>et al.</i> (2023) Induction of humoral and cell-mediated immunity in mice by chitosan-curdlan composite nanoparticles administered intranasally and subcutaneously <u>J Drug Deliv Sci Technol. 86: 104704.</u> Zhang, S. <i>et al.</i> (2023) Select symbionts drive high IgA levels in the mouse intestine. <u>Cell Host Microbe. 31 (10): 1620-1638.e7.</u> Amaral, R. <i>et al.</i> (2025) Chitosan Nanoparticles for Enhanced Immune Response and Delivery of Multi-Epitope <i>Helicobacter pylori</i> Vaccines in a BALB/c Mouse Model. <u>Pharmaceutics. 17 (1): 132.</u> 			
Storage Guarantee	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.			
Health And Safety	Material Safety Datasheet documentation #10077 available at: <u>https://www.bio-rad-antibodies.com/SDS/STAR137</u> 10077			
Regulatory	For research purposes only			

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) FITC, HRP

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-r	Worldwide ad.com	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-r	Europe ad.com	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-ra	To ^{id.} mid a	
batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M428699:240301'							

Printed on 30 Jan 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint