

Datasheet: STAR137F BATCH NUMBER 148278

Description:	GOAT ANTI MOUSE IgA:FITC
Specificity:	lgA
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			Neat - 1/10
Immunohistology - Frozen	•			
Immunohistology - Paraffin			•	
Immunofluorescence	•			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Mouse			
Product Form	Purified IgG conju	ugated to Fluorescein Isotl	niocyanate Isomer 1 (FITC) - liquid
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)	
	FITC	490	525	

Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.1% Sodium Azide (NaN ₃)
Approx. Protein	laG concentration 0.5mg/ml

Concentrations

Immunogen	Mouse IgA paraproteins.
External Database	UniDuct
Links	UniProt:
	P01878 Related reagents
	Entrez Gene:
	16061 Igh-VJ558 Related reagents
RRID	AB_2075636
Specificity	Goat anti Mouse IgA antibody recognizes Mouse IgA and has been cross absorbed against mouse IgM, IgG1, IgG2a, IgG2b and IgG3.
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	 Capoccia, B.J. <i>et al.</i> (2011) Transcription factor MIST1 in terminal differentiation of mouse and human plasma cells. <u>Physiol Genomics. 43 (3): 174-86.</u> Yamashita, H. <i>et al</i> (2012) Overcoming food allergy through acquired tolerance conferred by transfer of Tregs in a murine model. <u>Allergy. 67: 201-9.</u>
	3. Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced
	3. Won, Y.S. et al. (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced
Storage	 3. Won, Y.S. et al. (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. <u>J Nat Med. 71 (1): 68-75.</u> 4. Ramstead, A.G. et al. (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88
Storage	 Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. <u>J Nat Med. 71 (1): 68-75.</u> Ramstead, A.G. <i>et al.</i> (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary <i>Coxiella burnetii</i> Infection. <u>Infect Immun. 84 (4): 940-9.</u>
Storage	 Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. <u>J Nat Med. 71 (1): 68-75.</u> Ramstead, A.G. <i>et al.</i> (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary <i>Coxiella burnetii</i> Infection. <u>Infect Immun. 84 (4): 940-9.</u> Store at +4°C or at -20°C if preferred.
Storage	 Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. <u>J Nat Med. 71 (1): 68-75.</u> Ramstead, A.G. <i>et al.</i> (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary <i>Coxiella burnetii</i> Infection. <u>Infect Immun. 84 (4): 940-9.</u> Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and
Storage	 3. Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. J Nat Med. 71 (1): 68-75. 4. Ramstead, A.G. <i>et al.</i> (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary <i>Coxiella burnetii</i> Infection. Infect Immun. 84 (4): 940-9. Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this
	 3. Won, Y.S. <i>et al.</i> (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. J Nat Med. 71 (1): 68-75. 4. Ramstead, A.G. <i>et al.</i> (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary <i>Coxiella burnetii</i> Infection. Infect Immun. 84 (4): 940-9. Store at +4°C or at -20°C if preferred. This product should be stored undiluted. Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

America

North & South Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint