

Datasheet: STAR137F

BATCH NUMBER 165122

Description:	GOAT ANTI MOUSE IgA:FITC
Specificity:	IgA
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
Immunofluorescence	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species Mouse

Product Form Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Antiserum Preparation Antisera to mouse IgA were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution Phosphate buffered saline

Preservative Stabilisers <0.1% Sodium Azide (NaN₃)

Approx. Protein IgG concentration 0.5mg/ml

Concentrations

Immunogen Mouse IgA paraproteins.

External Database Links

UniProt:

[P01878](#) [Related reagents](#)

Entrez Gene:

[16061](#) Igh-VJ558 [Related reagents](#)

RRID AB_2075636

Specificity **Goat anti Mouse IgA antibody** recognizes Mouse IgA and has been cross absorbed against mouse IgM, IgG1, IgG2a, IgG2b and IgG3.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Capoccia, B.J. *et al.* (2011) Transcription factor MIST1 in terminal differentiation of mouse and human plasma cells. [Physiol Genomics. 43 \(3\): 174-86.](#)
2. Yamashita, H. *et al.* (2012) Overcoming food allergy through acquired tolerance conferred by transfer of Tregs in a murine model. [Allergy. 67: 201-9.](#)
3. Won, Y.S. *et al.* (2017) Green tea cultivar 'Benifuuki' potentiates split vaccine-induced immunoglobulin A production. [J Nat Med. 71 \(1\): 68-75.](#)
4. Ramstead, A.G. *et al.* (2016) Roles of Toll-Like Receptor 2 (TLR2), TLR4, and MyD88 during Pulmonary *Coxiella burnetii* Infection. [Infect Immun. 84 \(4\): 940-9.](#)

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/STAR137F10040>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M387535:210629'

