

Datasheet: STAR145

BATCH NUMBER 158507

| | |
|----------------------|---------------------|
| Description: | GOAT ANTI HUMAN IgM |
| Specificity: | IgM |
| Format: | Purified |
| Product Type: | Polyclonal Antibody |
| Isotype: | Polyclonal IgG |
| Quantity: | 1 mg |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------------------|-----|----|----------------|--------------------|
| Flow Cytometry | | | ▪ | |
| Immunohistology - Frozen | | | ▪ | |
| Immunohistology - Paraffin | | | ▪ | |
| ELISA | ▪ | | | |
| Western Blotting | | | ▪ | |
| Immunofluorescence | | | ▪ | |

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

| | |
|---------------------------------------|---|
| Target Species | Human |
| Product Form | Purified IgG - liquid |
| Antiserum Preparation | Antisera to human IgM were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. |
| Buffer Solution | Borate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide |
| Approx. Protein Concentrations | IgG concentration 1.0 mg/ml |

Immunogen Human IgM paraproteins.

External Database

Links

UniProt:

[P01871](#) [Related reagents](#)

Entrez Gene:

[3507](#) IGHM [Related reagents](#)

RRID AB_1102677

Specificity **Goat anti Human IgM antibody** recognizes the heavy chain of human IgM. This antibody has been cross absorbed against human IgA and IgG. Goat anti Human IgM antibody might cross react with IgM from other species.

References

1. Sharp, C.P. *et al.* (2012) Virologic and clinical features of primary infection with human parvovirus 4 in subjects with hemophilia: frequent transmission by virally inactivated clotting factor concentrates. [Transfusion. 52: 1482-9.](#)
2. Whelan, S.F. *et al.* (2013) Distinct characteristics of antibody responses against factor VIII in healthy individuals and in different cohorts of hemophilia A patients. [Blood. 121: 1039-48.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10077 available at: <https://www.bio-rad-antibodies.com/SDS/STAR145>
10077

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Rabbit Anti Goat IgG (Fc) (STAR122...) [FITC](#), [HRP](#)

North & South America Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M384393:210513'

Printed on 01 Mar 2024

