

Datasheet: STAR145F

BATCH NUMBER 167760

Description:	GOAT ANTI HUMAN IgM:FITC
Specificity:	IgM
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			Neat - 1/10
Immunohistology - Frozen	▪			
Immunohistology - Paraffin			▪	
Immunofluorescence	▪			Neat - 1/10

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Human

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Antiserum Preparation

Antisera to human IgM were raised by repeated immunisation of goats with purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

<0.1% Sodium Azide (NaN₃)

Approx. Protein Concentrations

IgG concentration 1.0 mg/ml

Immunogen Human IgM paraproteins.

External Database Links

UniProt:
[P01871](#) [Related reagents](#)

Entrez Gene:
[3507](#) IGHM [Related reagents](#)

RRID AB_1102678

Specificity **Goat anti Human IgM antibody** recognizes the heavy chain of human IgM. This antibody has been cross absorbed against human IgA and IgG. Goat anti Human IgM antibody might cross react with IgM from other species.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Sharp, C.P. *et al.* (2012) Virologic and clinical features of primary infection with human parvovirus 4 in subjects with hemophilia: frequent transmission by virally inactivated clotting factor concentrates. [Transfusion. 52: 1482-9.](#)
2. Whelan, S.F. *et al.* (2013) Distinct characteristics of antibody responses against factor VIII in healthy individuals and in different cohorts of hemophilia A patients. [Blood. 121: 1039-48.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/STAR145F>
10040

Regulatory For research purposes only

Related Products

Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Worldwide Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Europe Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

'M421288:230706'

Printed on 01 Mar 2024
