

## Datasheet: STAR147A

**BATCH NUMBER 166220**

<b>Description:</b>	GOAT ANTI HUMAN IgE:Alk. Phos.
<b>Specificity:</b>	IgE
<b>Format:</b>	Alk. Phos.
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	0.5 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			▪	
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			1/1000 - 1/10000
Immunoprecipitation			▪	
Western Blotting	▪			1/1000 - 1/5000

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Reacts with: Rhesus Monkey

**N.B.** Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

#### Product Form

Purified IgG conjugated to Alkaline Phosphatase - liquid

#### Antiserum Preparation

Antiserum to human IgE was raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

#### Buffer Solution

50mM HEPES, 0.1M NaCl, 1mM MgCl<sub>2</sub>, 0.1mM ZnCl<sub>2</sub>

<b>Preservative</b>	0.09% Sodium Azide
<b>Stabilisers</b>	0.2% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 0.5 mg/ml.
<b>Immunogen</b>	Native human IgE.
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P01854</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">3497</a> IGHE    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_1221598
<b>Specificity</b>	<p><b>Goat anti Human IgE polyclonal antibody</b> recognizes human IgE. Minimal cross-reactivity (&lt;0.1%) is expected with other human immunoglobulins and light chains.</p> <p>Goat anti Human IgE in a biotinylated format has been successfully employed as a detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE.</p>
<b>References</b>	<ol style="list-style-type: none"> <li>1. Khan, F.M. <i>et al.</i> (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. <a href="#">Allergy Asthma Clin Immunol. 8 (1): 1.</a></li> <li>2. Pinot de Moira, A. <i>et al.</i> (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. <a href="#">Infect Immun. 81 (1): 23-32.</a></li> <li>3. Zheng, W. <i>et al.</i> (2016) Upregulated expression of substance P in basophils of the patients with chronic spontaneous urticaria: induction of histamine release and basophil accumulation by substance P. <a href="#">Cell Biol Toxicol. 32 (3): 217-28.</a></li> <li>4. Jin Y <i>et al.</i> (2015) Allergenic response to squid (<i>Todarodes pacificus</i>) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. <a href="#">Food Chem Toxicol. 76: 86-93.</a></li> <li>5. Wei, J.F. <i>et al.</i> (2009) Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. <a href="#">BMC Immunol. 10: 21.</a></li> <li>6. Tsai, C.T. <i>et al.</i> (2018) Isotype-specific agglutination-PCR (ISAP): A sensitive and multiplex method for measuring allergen-specific IgE. <a href="#">J Allergy Clin Immunol. 141 (5): 1901-4.e15.</a></li> </ol>
<b>Storage</b>	<p>Store at +4°C.  DO NOT FREEZE</p> <p>This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.</p>
<b>Guarantee</b>	12 months from date of despatch.

---

**Health And Safety Information**      Material Safety Datasheet documentation #10089 available at:  
<https://www.bio-rad-antibodies.com/SDS/STAR147A>  
10089

---

**Regulatory**                      For research purposes only.

---

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)  
'M421294:230706'

**Printed on 01 Mar 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)