Datasheet: STAR147B BATCH NUMBER 167612

Description:	GOAT ANTI HUMAN IgE:Biotin
Specificity:	IgE
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications	This product has been re derived from testing withi communications from the	n our labo	oratories, p	peer-reviewed publication	tions or personal	
	information. For general	•				
	rad-antibodies.com/proto	<u>cols</u> .				
		Yes	Νο	Not Determined	Suggested Dilution	
	Flow Cytometry			•		
	Immunohistology - Frozen			•		
	Immunohistology - Paraffin			•		
	ELISA				1:10000 - 1:100000	
	Western Blotting				1:10000 - 1:100000	
	Where this product has n	ot been te	ested for u	se in a particular tech	inique this does not	
	necessarily exclude its use in such procedures. Suggested working dilutions are given as					
	a guide only. It is recomn system using appropriate			•	or use in their own	
Target Species	Human					
Species Cross	Reacts with: Rhesus Mor	nkey				
Reactivity	N.B. Antibody reactivity a	and workir	ng conditio	ns may vary between	species. Cross	
	reactivity is derived from	testing wi	thin our la	poratories, peer-reviev	wed publications or	
	personal communications	-		•	•	
	further information.		g			
Product Form	Purified IgG conjugated t	o Biotin -	liquid			
Antiserum Preparation	n Antisera to human IgE we antigen. Purified IgG was agarose beads			•	e i i i	
Buffer Solution	Phosphate buffered salin	e.				

Preservative 0.09% Sodium Azide (NaN3) Stabiliers 0.2% Bovine Serum Albumin Approx. Protein Concentrations IgG concentration 1.0 mg/ml. Immunogen Native human IgE. External Database Links Entrez Gene: 3497 3497 IGHE Related reagents Entrez Gene: 3497 Gotta anti Human IgE polyclonal antibody recognizes human IgE. Minimal cross- reactivity (<0.1%) is expected with other human immunoglobulins and light chains. Goat anti Human IgE in a biotinylated format has been successfully employed as a detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE. References 1. Khan, F.M. <i>et al.</i> (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. Allergy Asthma Clin Immunol. 8 (1): 1. 2. Pinot de Moira, A. <i>et al.</i> (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and Immunity to reinfection in childhood schistosome and hokworm coinfections. Infect Immun. 81 (1): 23-32. 3. Zheng, W. <i>et al.</i> (2016) Upregulated expression of substance P in basophils of the patients with chronic spontanecus utuaria: induction of histamine release and basophil accumulation by substance P. Cell Biol Toxicol. 32 (3): 217-28. 4. Jin Y <i>et al.</i> (2016) Upregulated expression of substance P in basophils of the patients with chronic spontanecus utuaria: induction of histamine release and skin edems by N49 phospholipase A2. BMC Immunol. 10. 21. 5		
Concentrations igG concentration 1.0 mg/ml. Immunogen Native human IgE. External Database UniProt: P01854 P01854 Entrez Gene: 3497 IGHE Related reagents RRD AB_10672882 Specificity Goat anti Human IgE polycional antibody recognizes human IgE. Minimal cross- reactivity (<0.1%) is expected with other human immunoglobulins and light chains. Goat anti Human IgE in a biotinylated format has been successfully employed as a detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE. References 1. Khan, F.M. et al. (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroaliergen-specific Immunoglobulin-E. Allergy Asthma Clin Immunol. 8 (1): 1. 2. Pinot de Moira, A. et al. (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. Infect Immun. 81 (1): 23-22. 3. Zheng, W. et al. (2016) Allergenic response to squid (<i>Todardes pacificus</i>) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. Food Chem Toxicol. 76: 86-93. 5. Wei, J.F. et al. (2018) Isotype-specific IgE. J. Allergy Clin Immunol. 141 (5): 19014-e15. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.		
External Database UniProt: PD1854 Related reagents Entrez Gene: 3497 3497 IGHE RRID AB_10672882 Specificity Goat anti Human IgE polyclonal antibody recognizes human IgE. Minimal cross-reactivity (<0.1%) is expected with other human immunoglobulins and light chains. Goat anti Human IgE in a biotinylated format has been successfully employed as a detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE. References 1. Khan, F.M. <i>et al.</i> (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. Allergy Asthma Clin Immunol. 8 (1): 1. 2. Pinot de Moira, A. <i>et al.</i> (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. Infect Immun. 81 (1): 23-32. 3. Zheng, W. <i>et al.</i> (2015) Allergenic response to squid (<i>Todarodes pacificus</i>) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. Food Chem Toxicol. 76: 86-93. 5. Wei, J.F. <i>et al.</i> (2019) Induction of mast cell accumulation, histamine release and skin edem by N49 phospholipase A2. BMC Immunol. 10: 21. 6. Tsai, C.T. <i>et al.</i> (2018) Uspre-specific IgE. J Allergy Clin Immunol. 141 (5): 1901-4.e15. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted, Should this product contain a precipitate we recommend microcentrifugation before use. <th></th> <th>IgG concentration 1.0 mg/ml.</th>		IgG concentration 1.0 mg/ml.
Links UniProt: P01854 Related reagents Entrez Gene: 3497 3497 IGHE RRID AB_10672882 Specificity Goat anti Human IgE polyclonal antibody recognizes human IgE. Minimal cross-reactivity (<0.1%) is expected with other human immunoglobulins and light chains. Goat anti Human IgE in a biotinylated format has been successfully employed as a detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE. References 1. Khan, F.M. et al. (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. Allergy Asthma Clin Immunol. 8 (1): 1. 2. Pinot de Moira, A. et al. (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. Infect Immun. 81 (1): 23-32. 3. Zheng, W. et al. (2016) Upregulated expression of substance P in basophils of the patients with chronic spontaneous urticaria: induction of histamine release and basophil accumulation by substance P. Cell Biol Toxicol. 32 (3): 217-28. 4. Jin Y et al. (2016) Mergenic response to squid (Todarades pacificus) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. Eood Chem Toxicol. 76: 86-93. 5. Wei, J.F. et al. (2009) Induction of mast cell accumulation, histamine release and skin edem by N49 phospholipase A2. <u>BMC Immunol. 10: 21.</u> 6. Tsai, C.T. et al. (2010) Isotype-specific aggluination-PCR (ISAP	Immunogen	Native human IgE.
RRID AB_10672882 Specificity Goat anti Human IgE polyclonal antibody recognizes human IgE. Minimal cross-reactivity (<0.1%) is expected with other human immunoglobulins and light chains.		P01854 Related reagents Entrez Gene:
Specificity Goat anti Human IgE polyclonal antibody recognizes human IgE. Minimal cross-reactivity (<0.1%) is expected with other human immunoglobulins and light chains.		
reactivity (<0.1%) is expected with other human immunoglobulins and light chains. Goat anti Human IgE in a biotinylated format has been successfully employed as a detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE. References 1. Khan, F.M. <i>et al.</i> (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. <u>Alleray Asthma Clin Immunol. 8 (1): 1.</u> 2. Pinot de Moira, A. <i>et al.</i> (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. <u>Infect Immun. 81 (1): 23-32.</u> 3. Zheng, W. <i>et al.</i> (2016) Upregulated expression of substance P in basophils of the patients with chronic spontaneous urticaria: induction of histamine release and basophil accumulation by substance P. <u>Cell Biol Toxicol. 32 (3): 217-28.</u> 4. Jin Y <i>et al.</i> (2009) Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. <u>BMC Immunol. 10: 21.</u> 6. Tsai, C.T. <i>et al.</i> (2018) Isotype-specific IgE. J Allergy Clin Immunol. 141 (5): 1901-4.e15. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.	RRID	AB_10672882
detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE. References 1. Khan, F.M. et al. (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. Allergy Asthma Clin Immunol. 8 (1): 1. 2. Pinot de Moira, A. et al. (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. Infect Immun. 81 (1): 23-32. 3. Zheng, W. et al. (2016) Upregulated expression of substance P in basophils of the patients with chronic spontaneous urticaria: induction of histamine release and basophil accumulation by substance P. Cell Biol Toxicol. 32 (3): 217-28. 4. Jin Y et al. (2015) Allergenic response to squid (<i>Todarodes pacificus</i>) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. Food Chem Toxicol. 76: 86-93. 5. Wei, J.F. et al. (2009) Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. BMC Immunol. 10: 21. 6. Tsai, C.T. et al. (2018) Isotype-specific agglutination-PCR (ISAP): A sensitive and multiplex method for measuring allergen-specific IgE. J Allergy Clin Immunol. 141 (5): 1901-4.e15. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.	Specificity	
fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. Allergy Asthma Clin Immunol. 8 (1): 1. 2. Pinot de Moira, A. <i>et al.</i> (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. Infect Immun. 81 (1): 23-32. 3. Zheng, W. <i>et al.</i> (2016) Upregulated expression of substance P in basophils of the patients with chronic spontaneous urticaria: induction of histamine release and basophil accumulation by substance P. Cell Biol Toxicol. 32 (3): 217-28. 4. Jin Y <i>et al.</i> (2015) Allergenic response to squid (<i>Todarodes pacificus</i>) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. Food Chem Toxicol. 76: 86-93. 5. Wei, J.F. <i>et al.</i> (2009) Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. BMC Immunol. 10: 21. 6. Tsai, C.T. <i>et al.</i> (2018) Isotype-specific agglutination-PCR (ISAP): A sensitive and multiplex method for measuring allergen-specific IgE. J Allergy Clin Immunol. 141 (5): 1901-4.e15. Storage Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.		detection reagent in the development of a sensitive sandwich ELISA for the measurement
This product should be stored undiluted. Should this product contain a precipitate we recommend microcentrifugation before use.	References	 fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. <u>Allergy</u> <u>Asthma Clin Immunol. 8 (1): 1.</u> 2. Pinot de Moira, A. <i>et al.</i> (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. <u>Infect Immun. 81 (1): 23-32.</u> 3. Zheng, W. <i>et al.</i> (2016) Upregulated expression of substance P in basophils of the patients with chronic spontaneous urticaria: induction of histamine release and basophil accumulation by substance P. <u>Cell Biol Toxicol. 32 (3): 217-28.</u> 4. Jin Y <i>et al.</i> (2015) Allergenic response to squid (<i>Todarodes pacificus</i>) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. <u>Food Chem Toxicol. 76: 86-93.</u> 5. Wei, J.F. <i>et al.</i> (2009) Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. <u>BMC Immunol. 10: 21.</u> 6. Tsai, C.T. <i>et al.</i> (2018) Isotype-specific agglutination-PCR (ISAP): A sensitive and multiplex method for measuring allergen-specific IgE. <u>J Allergy Clin Immunol. 141 (5)</u>;
Guarantee 12 months from date of despatch.	Storage	This product should be stored undiluted. Should this product contain a precipitate we
	Guarantee	12 months from date of despatch.

10041		<u>itibodies.com/SDS/STAF</u>	<u> 147 D</u>	
For resea	rch purpose:	s only.		
+1 800 265 7376 : +1 919 878 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50
ail: antibody_sales_us@bio-r	ad.com		-rad.com	Email: antibody_sales_de@bio-rad.com
	+1 800 265 7376 : +1 919 878 3751 ail: antibody_sales_us@bio-ra	+1 800 265 7376 Worldwide : +1 919 878 3751 ail: antibody_sales_us@bio-rad.com	: +1 919 878 3751 Fax: +44 (0)1865 852 739 ail: antibody_sales_us@bio-rad.com Email: antibody_sales_uk@bio	+1 800 265 7376 Worldwide Tel: +44 (0)1865 852 700 Europe : +1 919 878 3751 Fax: +44 (0)1865 852 739

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint