Datasheet: STAR147F BATCH NUMBER 160990

Description:	GOAT ANTI HUMAN IgE:FITC
Specificity:	IgE
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications	This product has been derived from testing wi communications from t information. For genera rad-antibodies.com/pro	thin our laborate he originators. I al protocol recor	ories, pe Please i	eer-reviewed publication refer to references in	tions or personal dicated for further
			No	Not Determined	Suggested Dilution
	Flow Cytometry	•			
	Immunohistology - Frozer	n			
	Immunohistology - Paraff	in		•	
	a guide only. It is recor system using the appro	use in such pro nmended that th	ocedure he user	s. Suggested working titrates the antibody	g dilutions are given as
Target Species	Human				
Species Cross	Reacts with: Rhesus M	lonkey			
Reactivity	N.B. Antibody reactivity reactivity is derived from personal communication further information.	m testing within	our lab	oratories, peer-revie	wed publications or
Product Form	Purified IgG conjugated	d to Fluoresceir	n Isothio	cyanate Isomer 1 (F	ITC) - liquid
Max Ex/Em	Fluorophore	Excitation Max	(nm) E	Emission Max (nm)	
	FITC	490		525	
Preparation	Purified IgG prepared b	by affinity chrom	natograp	bhy	

Antiserum Preparation Antisera to human IgE were raised by repeated immunisations of goats with highly purified antigen.

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide 0.2% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Native human IgE.
External Database Links	UniProt: <u>P01854</u> <u>Related reagents</u> Entrez Gene: <u>3497</u> IGHE <u>Related reagents</u>
RRID	AB_1221599
Specificity	Goat anti Human IgE polyclonal antibody recognizes human IgE. Minimal cross-reactivity (<0.1%) is expected with other human immunoglobulins and light chains.
	Goat anti Human IgE in a biotinylated format has been successfully employed as a detection reagent in the development of a sensitive sandwich ELISA for the measurement of human IgE.
Flow Cytometry	Use 10 ul of the suggested working dilution to label 1×10^6 cells in 100ul.
References	 Khan, F.M. <i>et al.</i> (2012) Basophil activation test compared to skin prick test and fluorescence enzyme immunoassay for aeroallergen-specific Immunoglobulin-E. <u>Allergy</u> <u>Asthma Clin Immunol. 8 (1): 1.</u> Pinot de Moira, A. <i>et al.</i> (2013) Effects of treatment on IgE responses against parasite allergen-like proteins and immunity to reinfection in childhood schistosome and hookworm coinfections. <u>Infect Immun. 81 (1): 23-32.</u> Zheng, W. <i>et al.</i> (2016) Upregulated expression of substance P in basophils of the patients with chronic spontaneous urticaria: induction of histamine release and basophil accumulation by substance P. <u>Cell Biol Toxicol. 32 (3): 217-28.</u> Jin Y <i>et al.</i> (2015) Allergenic response to squid (<i>Todarodes pacificus</i>) tropomyosin Tod p1 structure modifications induced by high hydrostatic pressure. <u>Food Chem Toxicol. 76: 86-93.</u> Wei, J.F. <i>et al.</i> (2009) Induction of mast cell accumulation, histamine release and skin edema by N49 phospholipase A2. <u>BMC Immunol. 10: 21.</u>
Storage	Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarante	e	12 monuns	from date d	of despatch		
Health Ar Informatio	•	https://www	•	heet documentation #10 htibodies.com/SDS/STA		at:
		10041				
		_		a anh		
Regulator Related	ry d Product	For researc	:h purpose:	soniy		
Related Recomm	-	S sful Reager BUF070A)		sony		
Related Recomm HUMAN SI HUMAN SI	d Product	S eful Reager BUF070A) BUF070B) 376		Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint