

Datasheet: STAR3010F

BATCH NUMBER 165339

Description:	GOAT ANTI RAT IgG/IgM (H/L):FITC (MOUSE ADSORBED)
Specificity:	IgG IgM (H/L)
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
FLISA	▪			1/100 - 1/400

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species

Rat

Species Cross Reactivity

Does not react with: Mouse

Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Preparation

Purified IgG prepared by affinity chromatography on pooled rat IgG linked to agarose

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers

<0.1% Sodium Azide (NaN₃)

Approx. Protein

IgG concentration 1.0 mg/ml

Concentrations

Immunogen Purified rat IgM and IgG

Specificity **Goat anti Rat IgG/IgM (H/L) polyclonal antibody** reacts with the heavy and light chains of rat IgM and IgG and show minimal cross reactivity with mouse immunoglobulins. This antibody has been cross-adsorbed against pooled mouse sera and purified mouse paraproteins.

References

1. Westerweel, P.E. *et al.* (2010) Modulation of TGF- β /BMP-6 expression and increased levels of circulating smooth muscle progenitor cells in a type I diabetes mouse model. [Cardiovasc Diabetol. 9: 55.](#)
2. Iver, S. *et al.* (2019) Probing BAK and BAX Activation and Pore Assembly with Cytochrome c Release, Limited Proteolysis, and Oxidant-Induced Linkage. [Methods Mol Biol. 1877: 201-16.](#)

Storage This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/STAR3010F10040>

Regulatory For research purposes only

North & South America Tel: +1 800 265 7376
Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700
Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21
Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M401963:220718'

Printed on 18 Jan 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)