

Datasheet: STAR4010B

BATCH NUMBER 169207

Description:	GOAT ANTI RABBIT IgG/IgM:Biotin (HUMAN ADSORBED)
Specificity:	IgG IgM
Format:	Biotin
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	•			
Immunohistology - Paraffin	•			
ELISA	•			1/5000 - 1/20,000
Western Blotting	•			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Rabbit
Species Cross Reactivity	Does not react with:Human
Product Form	Purified IgG conjugated to Biotin - liquid

Antiserum Preparation Antisera to rabbit IgG and IgM were raised by repeated immunizations of goats with normal rabbit IgG and IgM.

The Ig fraction was purified by affinity chromatography on pooled rabbit IgG and IgM

covalently linked to agarose.

The preparation has been cross absorbed with pooled human sera and purified human paraproteins to remove any potential cross reactivity.

Buffer Solution	Phosphate buffered saline

Preservative Stabilisers	<0.1% Sodium Azide (NaN ₃)		
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml		
Immunogen	Normal rabbit IgG + IgM.		
External Database Links	UniProt:		
	P01780 Related reagents		
	P03988 Related reagents		
	P04221 Related reagents		
Specificity	Goat anti Rabbit IgG/IgM antibody reacts with the heavy and light chains of rabbit IgG and IgM and with the light chains of rabbit IgA as demonstrated by ELISA and flow cytometry.		
References	 Iver, S. et al. (2019) Probing BAK and BAX Activation and Pore Assembly with Cytochrome c Release, Limited Proteolysis, and Oxidant-Induced Linkage. Methods Mol Biol. 1877: 201-16. Campos-Ordoñez, T. et al. (2021) Chronic exposure to cyclohexane induces stereotypic circling, hyperlocomotion, and anxiety-like behavior associated with atypical c-Fos expression in motor- and anxiety-related brain regions. Behav Brain Res. Nov 12;113664. 		
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.		
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.		
Guarantee	Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.		
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/STAR4010B 10040		

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M428745:240301'

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint