

Datasheet: STAR4010P

BATCH NUMBER 167318

Description:	GOAT ANTI RABBIT IgG/IgM:HRP (HUMAN ADSORBED)
Specificity:	IgG IgM
Format:	HRP
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunohistology - Frozen	▪			
Immunohistology - Paraffin	▪			
ELISA	▪			1/4000 - 1/8000

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

Target Species	Rabbit
Species Cross Reactivity	Does not react with:Human
Product Form	Purified IgG conjugated to Horseradish Peroxidase (HRP) - liquid

Antiserum Preparation Antisera to rabbit IgG and IgM were raised by repeated immunizations of goats with normal rabbit IgG and IgM.
The Ig fraction was purified by affinity chromatography on pooled rabbit IgG and IgM covalently linked to agarose.
The preparation has been cross absorbed with pooled human sera and purified human paraproteins to remove any potential cross reactivity.

Buffer Solution	Phosphate buffered saline
Preservative	50% glycerol

Stabilisers

Immunogen Normal rabbit IgG + IgM.

External Database Links

UniProt:

[P01780](#) [Related reagents](#)
[P03988](#) [Related reagents](#)
[P04221](#) [Related reagents](#)

Specificity **Goat anti Rabbit IgG/IgM antibody** reacts with the heavy and light chains of rabbit IgG and IgM and with the light chains of rabbit IgA as demonstrated by ELISA and flow cytometry.

References

1. Iver, S. *et al.* (2019) Probing BAK and BAX Activation and Pore Assembly with Cytochrome c Release, Limited Proteolysis, and Oxidant-Induced Linkage. [Methods Mol Biol. 1877: 201-16.](#)
2. Campos-Ordoñez, T. *et al.* (2021) Chronic exposure to cyclohexane induces stereotypic circling, hyperlocomotion, and anxiety-like behavior associated with atypical c-Fos expression in motor- and anxiety-related brain regions. [Behav Brain Res. Nov 12;113664.](#)

Storage Store at -20°C only.
Storage in frost-free freezers is not recommended.
This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee Guaranteed for 12 months from the date of despatch or until the date of expiry, whichever comes first. Please see label for expiry date.

Health And Safety Information Material Safety Datasheet documentation #10106 available at: <https://www.bio-rad-antibodies.com/SDS/STAR4010P10106>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M421339:230706'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)