

Datasheet: STAR69 BATCH NUMBER 159653

Description:	GOAT F(ab')2 ANTI RAT IgG:FITC (MOUSE ADSORBED)				
Specificity:	IgG				
Format:	FITC				
Product Type:	Polyclonal Antibody				
Isotype:	Polyclonal IgG				
Quantity:	0.5 ml				

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	information. For general protocol recommendations, please visit <u>www.bio-</u>						
	<u>rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			1/25 - 1/100		
	Immunohistology - Froze	n					
	Immunohistology - Paraff	fin		•			
	Where this antibody ha	as not been te	sted for	use in a particular tec	hnique this does not		
	necessarily exclude its	s use in such p	rocedur	es. Suggested working	g dilutions are given as		
	a guide only. It is recor	mmended that	the use	r titrates the antibody	for use in their own		
	system using appropria	ate negative/p	ositive c	ontrols.			
Target Species	Rat						
Product Form	F(ab')2 fragment of Ig0	G conjugated t	o Fluore	scein Isothiocyanate I	somer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Ma	x (nm)	Emission Max (nm)			
	FITC	490		525			
Antiserum Preparation	n Antisera to rat IgG wer antigen. Purified IgG w	•	•	-	vith highly purified		
Buffer Solution	Phosphate buffered sa	lline					
Preservative	0.09% Sodium Azide						
Stabilisers	0.5% Bovine Serum A	Albumin					
Approx. Protein Concentrations	F(ab')2 concentration (0.7 mg/ml					

Immunogen	Rat IgG.						
External Database Links	UniProt:P20759Related reagentsP20761Related reagentsP20762Related reagentsP20760Related reagents						
	Entrez Gene:						
	299354 Ighg Related reagents						
	362795 LOC362795 Related reagents						
	679045 LOC679045 Related reagents						
RRID	AB_321934						
Specificity	FITC conjugated Goat (F(ab') ₂ anti Rat IgG antibody recognises rat IgG. Cross- reactivity with mouse IgG has been reduced by adsorption, however, we recommend dilution in buffer containing 10% (v/v) normal mouse serum for use upon mouse tissues.						
Flow Cytometry	Use 10ul of the suggested working dilution to label 10 ⁶ cells in 100ul.						
References	 Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul. 1. Fowles, L. <i>et al.</i> (2000) Regulation of urokinase plasminogen activator gene transcription in the RAW264 murine macrophage cell line by macrophage colony-stimulating factor (CSF-1) is dependent upon the level of cell surface receptor. <u>Biochem</u>, J. 347: 313 - 320. 2. Modjtahedi, H. <i>et al.</i> (2012) Immunohistochemical discrimination of wild-type EGFR from EGFRVIII in fixed tumour specimens using anti-EGFR mAbs ICR9 and ICR10. <u>Br J</u> <u>Cancer. 106: 883-888</u> 3. Haraoka, M. <i>et al.</i> (1999) Neutrophil recruitment and resistance to urinary tract infection. <u>J Infect Dis. 180: 1220-9.</u> 4. Hang, L. <i>et al.</i> (2000) Interleukin-8 receptor knockout mice have subepithelial neutrophil entrapment and renal scarring following acute pyelonephritis. <u>J Infect Dis. 182: 1738-48.</u> 5. Franke, H. <i>et al.</i> (2005) P2X(7) receptor-mRNA and -protein in the mouse retina; changes during retinal degeneration in BALBCrds mice. <u>Neurochem Int. 47: 235-42.</u> 6. Parish, N.M. <i>et al.</i> (1999) Anti-CD44 treatment does not prevent the extravasation of autopathogenic T cells to the thyroid in experimental autoimmune thyroiditis. <u>Immunology. 97: 533-9.</u> 7. Song, L. <i>et al.</i> (2009) Lung vascular endothelial growth factor expression induces local myeloid dendritic cell activation. <u>Clin Immunol. 132: 371-84.</u> 9. Englezou, P.C. <i>et al.</i> (2015) P2X7R activation drives distinct IL-1 responses in dendritic cells compared to macrophages. <u>Cytokine. 74 (2): 293-304.</u> 						
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for						

		short term use (up to 4 weeks) and store the remaining aliquots at -20°C.						
		•	e antibody. Storage in sensitive and should be					
Guarante	96	12 months from date of						
Health A Informati	nd Safety ion	- t:						
Regulatory		For research purposes						
North & South America	Tel: +1 800 265 73 Fax: +1 919 878 3	3751	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739	·	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50			
To find a b		sales_us@bio-rad.com	Email: antibody_sales_uk@bio ict, please use our online s 'M385284:210513'		Email: antibody_sales_de@bio-rad.com			
Printed on 05 May 2025								

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint