

Datasheet: STAR69

Description:	GOAT F(ab') ₂ ANTI RAT IgG:FITC (MOUSE ADSORBED)
Specificity:	IgG
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.5 ml

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/100
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Rat		
Product Form	F(ab') ₂ fragment of IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Antiserum Preparation Antisera to rat IgG were raised by repeated immunisation of goat with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	0.5% Bovine Serum Albumin
Approx. Protein Concentrations	F(ab') ₂ concentration 0.7 mg/ml

Immunogen Rat IgG.

External Database

Links

UniProt:

[P20759](#) [Related reagents](#)
[P20761](#) [Related reagents](#)
[P20762](#) [Related reagents](#)
[P20760](#) [Related reagents](#)

Entrez Gene:

[299354](#) Ighg [Related reagents](#)
[362795](#) LOC362795 [Related reagents](#)
[679045](#) LOC679045 [Related reagents](#)

RRID AB_321934

Specificity **FITC conjugated Goat (F(ab')₂ anti Rat IgG antibody** recognises rat IgG. Cross-reactivity with mouse IgG has been reduced by adsorption, however, we recommend dilution in buffer containing 10% (v/v) normal mouse serum for use upon mouse tissues.

Flow Cytometry Use 10ul of the suggested working dilution to label 10⁶ cells in 100ul.

References

1. Fowles, L. *et al.* (2000) Regulation of urokinase plasminogen activator gene transcription in the RAW264 murine macrophage cell line by macrophage colony-stimulating factor (CSF-1) is dependent upon the level of cell surface receptor. [Biochem. J. 347: 313 - 320.](#)
 2. Modjtahedi, H. *et al.* (2012) Immunohistochemical discrimination of wild-type EGFR from EGFRvIII in fixed tumour specimens using anti-EGFR mAbs ICR9 and ICR10. [Br J Cancer. 106: 883-888](#)
 3. Haraoka, M. *et al.* (1999) Neutrophil recruitment and resistance to urinary tract infection. [J Infect Dis. 180: 1220-9.](#)
 4. Hang, L. *et al.* (2000) Interleukin-8 receptor knockout mice have subepithelial neutrophil entrapment and renal scarring following acute pyelonephritis. [J Infect Dis. 182: 1738-48.](#)
 5. Franke, H. *et al.* (2005) P2X(7) receptor-mRNA and -protein in the mouse retina; changes during retinal degeneration in BALBCrds mice. [Neurochem Int. 47: 235-42.](#)
 6. Parish, N.M. *et al.* (1999) Anti-CD44 treatment does not prevent the extravasation of autopathogenic T cells to the thyroid in experimental autoimmune thyroiditis. [Immunology. 97: 533-9.](#)
 7. Song, L. *et al.* (2011) Deletion of the murine scavenger receptor CD68. [J Lipid Res. 52: 1542-50.](#)
 8. Chapoval, S.P. *et al.* (2009) Lung vascular endothelial growth factor expression induces local myeloid dendritic cell activation. [Clin Immunol. 132: 371-84.](#)
 9. Englezou, P.C. *et al.* (2015) P2X7R activation drives distinct IL-1 responses in dendritic cells compared to macrophages. [Cytokine. 74 \(2\): 293-304.](#)
-

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee 12 months from date of despatch

Health And Safety Information Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M385284:210513'

Printed on 21 Mar 2022

© 2022 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)