

Datasheet: STAR71D800GA

BATCH NUMBER 165968

Description:	GOAT ANTI RAT IgG:DyLight®800 (MOUSE ADSORBED)
Specificity:	IgG (MOUSE ADSORBED)
Format:	DyLight®800
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			
Western Blotting	•			1/10000 - 1/50000
Immunofluorescence	-			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls

Target Species	Rat		
Product Form	Purified IgG conjugated to DyLight®800 - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	Dylight®800	777	794

Antiserum Preparation Antisera to rat IgG were raised by repeated immunisation of goats with highly purified antigen. Purified IgG prepared by affinity chromatography

Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	IgG concentration 1.0mg/ml

Immunogen	Rat IgG.			
External Database Links	UniProt: P20759 Related reagents P20762 Related reagents P20761 Related reagents P20760 Related reagents Entrez Gene: 299354 Ighg Related reagents 362795 LOC362795 Related reagents			
	679045 LOC679045 Related reagents			
RRID	AB_10852249			
Specificity	Goat anti Rat IgG (Mouse Adsorbed) antibody recognizes rat IgG. Cross-reactivity with mouse IgG has been minimised by adsorption.			
Flow Cytometry	Use 50ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul			
References	 Yang, X. et al. (2010) The role of the JAK2-STAT3 pathway in pro-inflammatory responses of EMF-stimulated N9 microglial cells. J Neuroinflammation. 7: 54. Tamayo, J. et al. (2001) Chemical sensors and biosensors in liquid environment based on microcantilevers with amplified quality factor. Ultramicroscopy. 86: 167-73. Pérez-Bosque A et al. (2004) Dietary plasma protein affects the immune response of weaned rats challenged with S. aureus Superantigen B. J Nutr. 134: 2667-72. Balan, P. et al. (2010) Immunomodulatory effects of ovine serum immunoglobulin in the growing rat. Animal. 4: 1702-8. Tulinská, J. et al. (2018) Humoral and cellular immune response in Wistar Han RCC rats fed two genetically modified maize MON810 varieties for 90 days (EU 7th Framework Programme project GRACE). Arch Toxicol. 92 (7): 2385-99. 			
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.			
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.			
Guarantee	12 months from date of despatch			
Acknowledgements	DyLight [®] is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries			
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/STAR71D800GA 10040			

Regulatory For research purposes only

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385536:210513'

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint