

## Datasheet: STAR74D488GA BATCH NUMBER 162268

Description:	GOAT ANTI MOUSE IgG:DyLight®488 (RAT ADSORBED)
Specificity:	IgG (RAT ADSORBED)
Format:	DyLight®488
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further						
	information. For general protocol recommendations, please visit <u>www.bio-</u>						
	rad-antibodies.com/protocols.						
		Yes	No	Not Determined	Suggested Dilution		
	Flow Cytometry	-			1/100 - 1/200		
	Immunofluorescence	•					
	Where this antibody has not been tested for use in a particular technique this does r						
	necessarily exclude its a guide only. It is reco system using appropri	mmended th	at the use	er titrates the antibody	ng dilutions are given as y for use in their own		
Target Species	Mouse						
Product Form	Purified IgG conjugate	ed to DyLight	®488 - liq	uid			
Max Ex/Em	Fluorophore	Excitation M	/lax (nm)	Emission Max (nm)			
	Dylight®488	493		518			
Antiserum Preparatio	<b>n</b> Antisera to mouse IgG antigen. Purified IgG p		• •		goats with highly purified		
Buffer Solution	Phosphate buffered saline						
Preservative Stabilisers	0.09% Sodium Azide						
Approx. Protein Concentrations	IgG concentration 1.0r						

Immunogen	Mouse IgG.				
External Database	U. D. t				
Links	UniProt:	Deletert			
			<u>reagents</u>		
			reagents		
	<u>P01863</u>	Related	reagents		
	Entrez Gene:				
	<u>16017</u> Igh	ng1	Related reagents		
	<u>380795</u> Al3	324046	Related reagents		
	<u>16016</u> Igh	ng2b	Related reagents		
	<u>16017</u> Igh	ng1	Related reagents		
	<u>380793</u> Igh-	า-1a	Related reagents		
	<u>380793</u> Igh	n-1a	Related reagents		
Synonyms	lgh-4				
RRID	AB_10845281				
Specificity	<b>Goat anti Mouse IgG (Rat Adsorbed) antibody</b> recognizes mouse IgG. Cross-reactivity with rat IgG has been minimised by adsorption.				
Flow Cytometry	Use 50ul of the suggested working dilution to label $1 \times 10^6$ cells in 100ul				
References	1. Jarrett, B.R. <i>et al.</i> (2010) <i>In vivo</i> mapping of vascular inflammation using multimodal imaging. <u>PLoS One. 5(10). pii: e13254.</u>				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.				
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.				
Guarantee	12 months from date of despatch				
Acknowledgements	DyLight <sup>®</sup> is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/STAR74D488GA 10040				
Regulatory	For research purposes only				

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-rad.c	com	Email: antibody_sales_uk@bio-rad	d.com	Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M385048:210513'

## Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint