

Datasheet: STAR74D550

BATCH NUMBER 159291

Description:	GOAT ANTI MOUSE IgG:DyLight®550 (RAT ADSORBED)
Specificity:	IgG (RAT ADSORBED)
Format:	DyLight®550
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-</u>				
	rad-antibodies.com/protocols.				
		Yes	No	Not Determined	Suggested Dilution
	Flow Cytometry	-			1/50 - 1/200
	Western Blotting	•			1/1000 - 1/10,000
	Immunofluorescence	•			1/50 - 1/200
	Where this antibody ha	as not been t	ested for u	use in a particular t	echnique this does not
	necessarily exclude its a guide only. It is recor system using appropria	mmended that	at the user	titrates the antiboo	king dilutions are given as dy for use in their own
Target Species	Mouse				
Product Form	Purified IgG conjugated to DyLight [®] 550 - liquid				
Max Ex/Em	Fluorophore	Excitation M	ax (nm)	Emission Max (nm)	
	Dylight®550	562		576	
Antiserum Preparation	Antisera to mouse IgG antigen. Purified IgG p				f goats with highly purified
Buffer Solution	Phosphate buffered saline				
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)				
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml				

Immunogen	Mouse IgG.				
External Database Links	UniProt:P01869Related reagentsP03987Related reagentsP01867Related reagentsP01864Related reagentsP01863Related reagentsP01863Related reagentsP01863Related reagentsP01863Related reagentsD0161Related reagents16017Ighg1Related reagents380795Al324046Related reagents16016Ighg2bRelated reagents16017Ighg1Related reagents16017Ighg1Related reagents16017Ighg1Related reagents16017Ighg1Related reagents380793Igh-1aRelated reagents380793Igh-1aRelated reagents				
Synonyms	Igh-4				
Specificity	Goat anti Mouse IgG (Rat Adsorbed) antibody recognizes mouse IgG. Cross-reactivity with rat IgG has been minimised by adsorption.				
Flow Cytometry	Use 50ul of the suggested working dilution to label 1×10^6 cells in 100ul.				
References	1. Jarrett, B.R. <i>et al.</i> (2010) <i>In vivo</i> mapping of vascular inflammation using multimodal imaging. <u>PLoS One. 5(10). pii: e13254.</u>				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C. Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended.				
Guarantee	12 months from date of despatch				
Acknowledgements	DyLight [®] is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.				
Health And Safety Information	Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/STAR74D550 10040				
Regulatory	For research purposes only				

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets

North & South Tel: +1 800 265 7376 America

Fax: +1 919 878 3751

Worldwide

Tel: +44 (0)1865 852 700 Europe Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com 'M398796:220624'

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

Printed on 01 Mar 2024

© 2024 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com