

Datasheet: STAR88D488GA

BATCH NUMBER 155632

Description:	DONKEY ANTI SHEEP/GOAT IgG:DyLight®488
Specificity:	IgG
Format:	DyLight®488
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	•			1/50 - 1/200
Immunofluorescence	-			1/50 - 1/200

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls

Sheep		
Reacts with: Goat		
reactivity is derive personal commun	ed from testing within our la ications from the originato	aboratories, peer-reviewed publication
Purified IgG conju	gated to DyLight [®] 488 - liq	uid
Fluorophore	Excitation Max (nm)	Emission Max (nm)
Dylight®488	493	518
	N.B. Antibody reareactivity is derived personal communiturther information. Purified IgG conjugations of the properties of the personal communiture of the personal commu	• ,

Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified IgG from a sheep serum pool.
RRID	AB_10848346
Specificity	Donkey anti Sheep/Goat IgG antibody recognizes both sheep and goat IgG, which are immunologically very similar and has been adsorbed against human, rabbit, mouse and rat serum to minimise cross-reactivity.
Flow Cytometry	Use 50ul of the suggested working dilution to label 1x10 ⁶ cells in 100ul.
References	1. Singh, M. et al. (1999) A recombinant measles virus expressing hepatitis B virus

- 1. Singh, M. *et al.* (1999) A recombinant measles virus expressing hepatitis B virus surface antigen induces humoral immune responses in genetically modified mice. <u>J Virol.</u> 73 (6): 4823-8.
- 2. Tedla, N. *et al.* (1998) Regulation of T lymphocyte trafficking into lymph nodes during an immune response by the chemokines macrophage inflammatory protein (MIP)-1 alpha and MIP-1 beta. J Immunol. 161 (10): 5663-72.
- 3. Turner J *et al.* (2002) *In vivo* IL-10 production reactivates chronic pulmonary tuberculosis in C57BL/6 mice. <u>J Immunol. 169 (11): 6343-51.</u>
- 4. Singh, S.K. *et al.* (2010) Melanin transfer in human skin cells is mediated by filopodia--a model for homotypic and heterotypic lysosome-related organelle transfer. <u>FASEB J. 24:</u> 3756-69.
- 5. Yekta, M.A. *et al.* (2011) Immunization of sheep with a combination of intiminγ, EspA and EspB decreases Escherichia coli O157:H7 shedding. <u>Vet Immunol Immunopathol. 140</u> (1-2): 42-6.
- 6. Guitton, C. *et al.* (2011) Protective cross talk between activated protein C and TNF signaling in vascular endothelial cells: implication of EPCR, noncanonical NF-{kappa}B, and ERK1/2 MAP kinases. Am J Physiol Cell Physiol. 300: C833-42.
- 7. Clinton, S.R. *et al.* (2010) Binding and activation of host plasminogen on the surface of *Francisella tularensis*. BMC Microbiol. 10: 76.
- 8. Chimote, A.A. *et al.* (2012) Disruption of kv1.3 channel forward vesicular trafficking by hypoxia in human T lymphocytes. <u>J Biol Chem. 287: 2055-67.</u>
- 9. Junqueira-Kipnis, A.P. *et al.* (2005) Interleukin-10 production by lung macrophages in CBA xid mutant mice infected with *Mycobacterium tuberculosis*. <u>Immunology</u>. 115: 246-52.
- 10. Nicol, M.Q. *et al.* (2012) A novel family of peptides with potent activity against influenza A viruses. J Gen Virol. 93: 980-6.
- 11. Singh B *et al.* (2015) *Moraxella catarrhalis* Binds Plasminogen To Evade Host Innate Immunity. Infect Immun. 83 (9): 3458-69.
- 12. Al-Jubair, T. *et al.* (2015) *Haemophilus influenzae* Type f Hijacks Vitronectin Using Protein H To Resist Host Innate Immunity and Adhere to Pulmonary Epithelial Cells. <u>J Immunol</u>. 195 (12): 5688-95.
- 13. McNeilly, T.N. *et al.* (2013) Suppression of ovine lymphocyte activation by Teladorsagia circumcincta larval excretory-secretory products. Vet Res. 44: 70.

14. Garza, J.J. et al. (2017) Serum-mediated Haemonchus contortus larval aggregation differs by larval stage and is enhanced by complement. Parasite Immunol. 39 (3)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation

before use

Guarantee 12 months from date of despatch

DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries. **Acknowledgements**

Health And Safety Material Safety Datasheet documentation #10040 available at: Information 10040: https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf

Regulatory For research purposes only

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M369797:200529'

Printed on 09 Feb 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint