

Datasheet: STAR88F

Description:	DONKEY ANTI SHEEP/GOAT IgG:FITC
Specificity:	IgG
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry			■	
Immunohistology - Frozen	■			1/100 - 1/200
Immunohistology - Paraffin			■	

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species	Sheep		
Species Cross Reactivity	Reacts with: Goat N.B. Antibody reactivity and working conditions may vary between species.		
Product Form	Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - liquid		
Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	FITC	490	525

Antiserum Preparation Antisera to sheep IgG were raised by repeated immunisation of donkeys with highly purified antigen. Purified IgG was prepared by affinity chromatography.

Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide
Stabilisers	1% Bovine Serum Albumin
Approx. Protein Concentrations	IgG concentration 1.0 mg/ml
Immunogen	Purified IgG from a sheep serum pool.

Specificity

Donkey anti Sheep/Goat IgG antibody recognizes both sheep and goat IgG, which are immunologically very similar and has been adsorbed against human, rabbit, mouse and rat serum to minimise cross-reactivity.

References

1. Singh, M. *et al.* (1999) A recombinant measles virus expressing hepatitis B virus surface antigen induces humoral immune responses in genetically modified mice. [J Virol. 73 \(6\): 4823-8.](#)
2. Tedla, N. *et al.* (1998) Regulation of T lymphocyte trafficking into lymph nodes during an immune response by the chemokines macrophage inflammatory protein (MIP)-1 alpha and MIP-1 beta. [J Immunol. 161 \(10\): 5663-72.](#)
3. Turner J *et al.* (2002) *In vivo* IL-10 production reactivates chronic pulmonary tuberculosis in C57BL/6 mice. [J Immunol. 169 \(11\): 6343-51.](#)
4. Singh, S.K. *et al.* (2010) Melanin transfer in human skin cells is mediated by filopodia--a model for homotypic and heterotypic lysosome-related organelle transfer. [FASEB J. 24: 3756-69.](#)
5. Yekta, M.A. *et al.* (2011) Immunization of sheep with a combination of intimin, EspA and EspB decreases *Escherichia coli* O157:H7 shedding. [Vet Immunol Immunopathol. 140 \(1-2\): 42-6.](#)
6. Guitton, C. *et al.* (2011) Protective cross talk between activated protein C and TNF signaling in vascular endothelial cells: implication of EPCR, noncanonical NF- κ B, and ERK1/2 MAP kinases. [Am J Physiol Cell Physiol. 300: C833-42.](#)
7. Clinton, S.R. *et al.* (2010) Binding and activation of host plasminogen on the surface of *Francisella tularensis*. [BMC Microbiol. 10: 76.](#)
8. Chimote, A.A. *et al.* (2012) Disruption of kv1.3 channel forward vesicular trafficking by hypoxia in human T lymphocytes. [J Biol Chem. 287: 2055-67.](#)
9. Junqueira-Kipnis, A.P. *et al.* (2005) Interleukin-10 production by lung macrophages in CBA xid mutant mice infected with *Mycobacterium tuberculosis*. [Immunology. 115: 246-52.](#)
10. Nicol, M.Q. *et al.* (2012) A novel family of peptides with potent activity against influenza A viruses. [J Gen Virol. 93: 980-6.](#)
11. Singh B *et al.* (2015) *Moraxella catarrhalis* Binds Plasminogen To Evade Host Innate Immunity. [Infect Immun. 83 \(9\): 3458-69.](#)
12. Al-Jubair, T. *et al.* (2015) *Haemophilus influenzae* Type f Hijacks Vitronectin Using Protein H To Resist Host Innate Immunity and Adhere to Pulmonary Epithelial Cells. [J Immunol. 195 \(12\): 5688-95.](#)
13. McNeilly, T.N. *et al.* (2013) Suppression of ovine lymphocyte activation by *Teladorsagia circumcincta* larval excretory-secretory products. [Vet Res. 44: 70.](#)
14. Garza, J.J. *et al.* (2017) Serum-mediated *Haemonchus contortus* larval aggregation differs by larval stage and is enhanced by complement. [Parasite Immunol. 39 \(3\)](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

Guarantee

Guaranteed until date of expiry. Please see product label.

Health And Safety Information

Material Safety Datasheet documentation #10041 available at:
10041: <https://www.bio-rad-antibodies.com/uploads/MSDS/10041.pdf>

**North & South
America**

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M356848:190827'

Printed on 11 Aug 2020

© 2020 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)