

Datasheet: STAR8D800GA

Description:	RABBIT F(ab') ₂ ANTI MOUSE IgG:DyLight®800
Specificity:	IgG
Format:	DyLight®800
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	0.1 mg

Product Details

RRID AB_10846935

Applications This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			
Western Blotting	▪			1/10000 - 1/50000
Immunofluorescence	▪			

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls

Target Species Mouse

Species Cross Reactivity Reacts with: Rat
N.B. Antibody reactivity and working conditions may vary between species.

Product Form Purified IgG conjugated to DyLight®800 - liquid

Max Ex/Em	Fluorophore	Excitation Max (nm)	Emission Max (nm)
	DyLight®800	777	794

Preparation Purified IgG was prepared from whole serum by affinity chromatography. F(ab')₂ fragments were prepared by pepsin digestion of the IgG followed by a gel filtration step to remove any remaining intact IgG or Fc fragments

Buffer Solution Phosphate buffered saline

Preservative Stabilisers 0.09% Sodium Azide

Approx. Protein Concentrations IgG concentration 1.0 mg/ml

Immunogen

Whole mouse IgG.

External Database Links**UniProt:**

[P01869](#) [Related reagents](#)
[P01865](#) [Related reagents](#)
[P03987](#) [Related reagents](#)
[P01867](#) [Related reagents](#)
[P01868](#) [Related reagents](#)
[P01864](#) [Related reagents](#)
[P01863](#) [Related reagents](#)

Entrez Gene:

[16017](#) Ighg1 [Related reagents](#)
[380793](#) Igh-1a [Related reagents](#)
[16016](#) Ighg2b [Related reagents](#)
[16017](#) Ighg1 [Related reagents](#)
[380793](#) Igh-1a [Related reagents](#)
[380795](#) AI324046 [Related reagents](#)
[380793](#) Igh-1a [Related reagents](#)

Synonyms

Igh-4

Specificity

Rabbit F(ab')₂ anti Mouse IgG antibody recognizes all subclasses of mouse IgG. Cross reactivity with rat IgG is expected. Cross reactivity with human serum proteins has been minimised by solid phase adsorption.

STAR8B is suitable as a bridge in a PAP / APAAP complex. Peroxidase anti-Peroxidase (PAP) complexes are formed by antibodies raised towards Horse Radish Peroxidase (HRP), binding to HRP molecules. Such complexes are used as signal amplification reagents in immunochemical staining techniques. It must be emphasised that the primary antibody must be raised in the same species as the antibodies used to form the PAP complex, so that a secondary "linker" antibody can be used to "bridge" the primary antibody to the PAP complex.

Flow CytometryUse 50ul of the suggested working dilution to label 1×10^6 cells in 100ul**References**

1. Seehafer, S.S. *et al.* (2011) Immunosuppression alters disease severity in juvenile Batten disease mice. [J Neuroimmunol. 230 \(1-2\): 169-72.](#)
2. Motallebzadeh, R. *et al.* (2012) Blocking lymphotoxin signaling abrogates the development of ectopic lymphoid tissue within cardiac allografts and inhibits effector antibody responses. [FASEB J. 26 \(1\): 51-62.](#)
3. Niven, J. *et al.* (2015) S100B Up-Regulates Macrophage Production of IL1 β and CCL22 and Influences Severity of Retinal Inflammation. [PLoS One. 10 \(7\): e0132688.](#)

Storage

Store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost free freezers is not recommended. This product is photosensitive and should be protected from light. Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use

Guarantee 18 months from date of despatch

Acknowledgements DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries

Health And Safety Information Material Safety Datasheet documentation #10040 available at:
10040: <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M353058:190409'

Printed on 20 May 2019

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)