

## Datasheet: STAR90F

**BATCH NUMBER 155146**

<b>Description:</b>	GOAT ANTI HUMAN IgG/A/M:FITC
<b>Specificity:</b>	IgG IgA IgM
<b>Format:</b>	FITC
<b>Product Type:</b>	Polyclonal Antibody
<b>Isotype:</b>	Polyclonal IgG
<b>Quantity:</b>	2 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			0.1 ug/ml - 0.5 ug/ml
Immunohistology - Frozen			▪	
Immunohistology - Paraffin			▪	
ELISA	▪			0.04 ug/ml - 0.2 ug/ml

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

#### Target Species

Human

#### Species Cross Reactivity

Does not react with: Mouse

#### Product Form

Purified IgG conjugated to Fluorescein Isothiocyanate Isomer 1 (FITC) - lyophilised

#### Reconstitution

Reconstitute with 1.0 ml distilled water

Care should be taken during reconstitution as the protein may appear as a film at the bottom of the vial. Bio-Rad recommend that the vial is gently mixed after reconstitution.

#### Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

#### Antiserum Preparation

Antisera to human Ig were raised by repeated immunisation of goats with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography.

<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative</b>	0.01% Sodium Azide
<b>Stabilisers</b>	1% Bovine Serum Albumin
<b>Approx. Protein Concentrations</b>	IgG concentration 2.0 mg/ml after reconstitution
<b>Immunogen</b>	Human IgG, IgA and IgM whole molecules
<b>External Database Links</b>	<p><b>UniProt:</b></p> <p><a href="#">P01859</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01860</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01857</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01861</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01876</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01877</a>    <a href="#">Related reagents</a></p> <p><a href="#">P01871</a>    <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b></p> <p><a href="#">3501</a> IGHG2    <a href="#">Related reagents</a></p> <p><a href="#">3502</a> IGHG3    <a href="#">Related reagents</a></p> <p><a href="#">3493</a> IGHA1    <a href="#">Related reagents</a></p> <p><a href="#">3494</a> IGHA2    <a href="#">Related reagents</a></p> <p><a href="#">3500</a> IGHG1    <a href="#">Related reagents</a></p> <p><a href="#">3503</a> IGHG4    <a href="#">Related reagents</a></p> <p><a href="#">3507</a> IGHM    <a href="#">Related reagents</a></p>
<b>RRID</b>	AB_567009
<b>Specificity</b>	<b>Goat anti Human IgG/A/M antibody</b> recognizes human Ig, binding to all major classes (IgG, IgA, IgM).
<b>Flow Cytometry</b>	Use 10ul of the suggested working dilution to label 10 <sup>6</sup> cells in 100ul.
<b>References</b>	<ol style="list-style-type: none"> <li>Lierz, M. <i>et al.</i> (2008) Avian Mycoplasma lipofaciens transmission to veterinarian. <a href="#">Emerg Infect Dis. 14: 1161-3.</a></li> <li>Vidyarthi, A. <i>et al.</i> (2017) Antibody response against PhoP efficiently discriminates among healthy individuals, tuberculosis patients and their contacts. <a href="#">PLoS One. 12 (3): e0173769.</a></li> <li>Jaki, T. <i>et al.</i> (2016) A false sense of security? Can tiered approach be trusted to accurately classify immunogenicity samples? <a href="#">J Pharm Biomed Anal. 128: 166-73.</a></li> </ol>
<b>Storage</b>	Prior to reconstitution store at +4°C. Following reconstitution store at +4°C or at -20°C if preferred.

This product should be stored undiluted.

Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Avoid repeated freezing and thawing as this may denature the antibody. Should this product contain a precipitate we recommend microcentrifugation before use.

---

<b>Guarantee</b>	12 months from date of despatch
------------------	---------------------------------

---

<b>Health And Safety Information</b>	Material Safety Datasheet documentation #10224 available at: <a href="https://www.bio-rad-antibodies.com/SDS/STAR90F10224">https://www.bio-rad-antibodies.com/SDS/STAR90F10224</a>
--------------------------------------	---

---

<b>Regulatory</b>	For research purposes only
-------------------	----------------------------

---

## Related Products

### Recommended Useful Reagents

[HUMAN SEROBLOCK \(BUF070A\)](#)

[HUMAN SEROBLOCK \(BUF070B\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://bio-rad-antibodies.com/datasheets)  
'M375760:210107'

**Printed on 13 Aug 2023**