

Datasheet: STAR9B

BATCH NUMBER 159655

Description:	RABBIT F(ab') ₂ ANTI MOUSE IgG:FITC
Specificity:	IgG
Format:	FITC
Product Type:	Polyclonal Antibody
Isotype:	Polyclonal IgG
Quantity:	1 mg

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Flow Cytometry	▪			1/25 - 1/100

Where this antibody has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the antibody for use in their own system using appropriate negative/positive controls.

Target Species

Mouse

Species Cross Reactivity

Reacts with: Rat

N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.

Product Form

F(ab')₂ fragment of IgG conjugated to Fluorescein Isothiocyanate Isomer I (FITC) - liquid

Max Ex/Em

Fluorophore	Excitation Max (nm)	Emission Max (nm)
FITC	490	525

Antiserum Preparation

Antisera to Mouse IgG were raised by repeated immunisation of rabbits with highly purified antigen. Purified IgG was prepared from whole serum by affinity chromatography. F(ab')₂ fragments were prepared by pepsin digestion of the IgG followed by a gel filtration step to remove the remaining intact IgG or Fc fragments.

Buffer Solution

Phosphate buffered saline

Preservative Stabilisers	0.09% Sodium Azide
Approx. Protein Concentrations	F(ab') ₂ concentration 1.0 mg/ml
Immunogen	Purified mouse IgG.
External Database Links	<p>UniProt:</p> <p>P01869 Related reagents</p> <p>P01865 Related reagents</p> <p>P03987 Related reagents</p> <p>P01864 Related reagents</p> <p>P01867 Related reagents</p> <p>P01868 Related reagents</p> <p>P01863 Related reagents</p> <p>Entrez Gene:</p> <p>16017 Ighg1 Related reagents</p> <p>380793 Igh-1a Related reagents</p> <p>16016 Ighg2b Related reagents</p> <p>16017 Ighg1 Related reagents</p> <p>380793 Igh-1a Related reagents</p> <p>380795 AI324046 Related reagents</p> <p>380793 Igh-1a Related reagents</p>
Synonyms	Igh-4
RRID	AB_321920
Specificity	<p>FITC conjugated Rabbit F(ab')₂ anti Mouse IgG antibody recognizes all subclasses of mouse IgG.</p> <p>Some cross reactivity with mouse IgM and IgA is expected, as is cross reactivity with rat IgG. Cross reactivity with human serum proteins has been minimised by solid phase adsorption.</p>
Flow Cytometry	Use 50ul of the suggested working dilution to label 10 ⁶ cells in 100ul.
References	<ol style="list-style-type: none"> O-charoenrat, P. <i>et al.</i> (2000) Epidermal growth factor-like ligands differentially up-regulate matrix metalloproteinase 9 in head and neck squamous carcinoma cells. Cancer Res. 60 (4): 1121-8. Lamote, I. <i>et al.</i> (2004) Influence of 17beta-estradiol, progesterone, and dexamethasone on diapedesis and viability of bovine blood polymorphonuclear leukocytes. J Dairy Sci. 87 (10): 3340-9. Dalli, J. <i>et al.</i> (2008) Annexin 1 mediates the rapid anti-inflammatory effects of

- neutrophil-derived microparticles. [Blood. 112 \(6\): 2512-9.](#)
4. Fleming, E.H. *et al.* (2006) Respiratory syncytial virus F envelope protein associates with lipid rafts without a requirement for other virus proteins. [J Virol. 80: 12160-70.](#)
 5. Peretti, M. *et al.* (2001) Expression of the three human major histocompatibility complex class II isotypes exhibits a differential dependence on the transcription factor RFXAP. [Mol Cell Biol. 21: 5699-709.](#)
 6. Krawczyk, M. *et al.* (2005) New functions of the major histocompatibility complex class II-specific transcription factor RFXANK revealed by a high-resolution mutagenesis study. [Mol Cell Biol. 25: 8607-18.](#)
 7. Frenzel, R. *et al.* (2006) The human thyrotropin receptor is predominantly internalized by beta-arrestin 2. [Endocrinology. 147: 3114-22.](#)
 8. Brancaleone, V. *et al.* (2011) Evidence for an anti-inflammatory loop centered on polymorphonuclear leukocyte formyl peptide receptor 2/lipoxin A4 receptor and operative in the inflamed microvasculature. [J Immunol. 186: 4905-14.](#)
 9. Waclavicek, M. *et al.* (2009) Analysis of the early response to TSST-1 reveals Vbeta-unrestricted extravasation, compartmentalization of the response, and unresponsiveness but not energy to TSST-1. [J Leukoc Biol. 85: 44-54.](#)
 10. Maderna, P. *et al.* (2010) FPR2/ALX receptor expression and internalization are critical for lipoxin A4 and annexin-derived peptide-stimulated phagocytosis. [FASEB J. 24: 4240-9.](#)
 11. Ioannou, N. *et al.* (2011) Anti-tumour activity of afatinib, an irreversible ErbB family blocker, in human pancreatic tumour cells. [Br J Cancer. 105: 1554-62.](#)
 12. Renshaw, D. *et al.* (2010) Downstream gene activation of the receptor ALX by the agonist annexin A1. [PLoS One. 5. pii: e12771.](#)
 13. Bena, S. *et al.* (2012) Annexin A1 interaction with the FPR2/ALX receptor: identification of distinct domains and downstream associated signaling. [J Biol Chem. 287: 24690-7.](#)
 14. Mehta, K. *et al.* (2016) Characterization of hepcidin response to holotransferrin in novel recombinant TfR1 HepG2 cells. [Blood Cells Mol Dis. 61: 37-45.](#)
 15. Puvanenthiran, S. *et al.* (2016) Impact of the putative cancer stem cell markers and growth factor receptor expression on the sensitivity of ovarian cancer cells to treatment with various forms of small molecule tyrosine kinase inhibitors and cytotoxic drugs. [Int J Oncol. 49 \(5\): 1825-38.](#)
 16. Ioannou, N. *et al.* (2013) Treatment with a combination of the ErbB (HER) family blocker afatinib and the IGF-IR inhibitor, NVP-AEW541 induces synergistic growth inhibition of human pancreatic cancer cells. [BMC Cancer. 13: 41.](#)
 17. Khan, T. *et al.* (2020) Synergistic activity of agents targeting growth factor receptors, CDKs and downstream signaling molecules in a panel of pancreatic cancer cell lines and the identification of antagonistic combinations: Implications for future clinical trials in pancreatic cancer [Oncology Reports. 44 \(6\): 2581-94.](#)
 18. Reitsma, L.M. *et al.* (2020) Effects of oral calcium bolus supplementation on intracellular polymorphonuclear leukocyte calcium levels and functionality in primiparous and multiparous dairy cows. [J Dairy Sci. 103 \(12\): 11876-88.](#)
 19. Nadkarni, S. *et al.* (2019) Identification of an activated neutrophil phenotype in polymyalgia rheumatica during steroid treatment: a potential involvement of immune cell cross-talk. [Clin Sci \(Lond\). 133 \(7\): 839-851.](#)
 20. da Silva, H.T. *et al.* (2021) Prophylactic use of an ultra-diluted complex on polymorphonuclear leukocyte function and respiratory scores of weaned Holstein calves

immediately after grouping [Res Soc Dev. 10 \(11\): e281101119582](#).

21. Lin, T.Y. *et al.* (2022) Radix Glycyrrhizae Preparata Induces Cell Cycle Arrest and Induced Caspase-Dependent Apoptosis in Glioblastoma Multiforme. [Neurol Int. 14 \(4\): 804-823](#).

22. Hung, C.M. *et al.* (2023) Sulforaphane-Induced Cell Mitotic Delay and Inhibited Cell Proliferation via Regulating CDK5R1 Upregulation in Breast Cancer Cell Lines. [Biomedicines. 11 \(4\): 996](#).

23. Al-Janaby, T. *et al.* (2024) The Combination of Afatinib With Dasatinib or Miransertib Results in Synergistic Growth Inhibition of Stomach Cancer Cells [World Journal of Oncology. 15 \(2\): 192-208](#).

24. Tsai, T.H. *et al.* (2021) RTA404, an Activator of Nrf2, Activates the Checkpoint Kinases and Induces Apoptosis through Intrinsic Apoptotic Pathway in Malignant Glioma. [J Clin Med. 10 \(21\): 4805](#).

Storage

This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.

Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. This product is photosensitive and should be protected from light.

Guarantee

12 months from date of despatch

Health And Safety Information

Material Safety Datasheet documentation #10040 available at: <https://www.bio-rad-antibodies.com/SDS/STAR9B>
10040

Regulatory

For research purposes only

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M385542:210513'

Printed on 19 Aug 2024

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)