Datasheet: TZA020 BATCH NUMBER 166617

Description:	ANTI MONO-ADP-RIBOSE
Specificity:	MONO-ADP-RIBOSE
Format:	Purified
Product Type:	Monoclonal Antibody
Clone:	AbD43647
Isotype:	Fab antibody
Quantity:	0.1 mg

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further							
	information. For general protocol recommendations, please visit <u>www.bio-</u> rad-antibodies.com/protocols.							
		Yes	No	Not Determined	Suggested Dilution			
	ELISA	-						
	Western Blotting	-						
	Immunofluorescence	-						
	Where this product has not been tested for use in a particular technique this does not							
	necessarily exclude its u	ise in sucł	n procedu	res. Suggested workin	g dilutions are given as			
	a guide only. It is recom	mended th	nat the us	er titrates the product f	or use in their own			
	system using appropriate	e negative	/positive	controls.				
Target Species	Protein/peptide tag							
Product Form	A monovalent human recombinant Fab selected from the HuCAL® phage display library, Expressed in a proprietary <i>E. coli.</i> strain. The antibody is tagged with a DYKDDDDK tag, a SpyTag version 2 (VPTIVMVDAYKRYK) and a His6-tag (HHHHHH) at the C-terminus of the antibody heavy chain. This antibody is supplied as a liquid.							
Preparation	Recombinant Fab antibody expressed in <i>E. coli</i> and purified by affinity chromatography							
Source	E.coli							
Buffer Solution	Phosphate buffered salir	ne						
Preservative Stabilisers	0.0095% MIT							

Carrier Free	Yes				
Approx. Protein Concentrations	Total protein concentration 0.5 mg/ml				
Immunogen	ARTKQTARKS(adpr)TGGKAPRKQLAGGK				
Specificity	Anti mono-ADP-ribose antibody, clone AbD43647, recognizes mono-adenosine diphosphate (ADP)-ribosylation. ADP-ribosylation is a reversible post-translational modification that occurs in multicellular organisms as well as some lower unicellular eukaryotes, but is absent in prokaryotes and yeast (<u>Bürkle 2005</u>). ADP ribosylation has been shown to play critical roles in many physiological and pathological processes, including bacterial pathogenesis and signaling and metabolism to control chromatin-related processes including transcription and DNA repair (<u>Bonfiglio et al. 2020</u> , <u>Bütepage et al. 2015</u>).				
	Members of the ADP-ribosyltransferase (ART) superfamily of proteins including the poly(ADP-ribose) polymerases (PARPs) subfamily, catalyze the transfer of ADP-ribose from nicotinamide adenine dinucleotide (NAD+) onto substrate protein via N-, O-, or S-glycosidic linkages. These transferases are able to transfer a single ADP-ribose residue to their substrate proteins, in a process known as mono-ADP-ribosylation (<u>Bütepage <i>et al.</i></u> 2015).				
	The anti mono-ADP-ribose antibody, clone AbD43647 is a mono-selective ADP-ribose antibody with a preference for mono-Ser-ADPr catalysed by the poly(ADP-ribose) polymerase 1 PARP1. Clone AbD43647 is tagged with a SpyTag2 at the C-terminus of the Fab heavy chain, enabling the user to couple it to a <u>SpyCatcher reagent</u> for conversion to alternative formats in less than an hour.				
	Anti mono-ADP-ribose antibody, clone AbD43647 is a mono-selective ADP-ribose antibody generated using an H3S10ADPr peptide as the antigen. This clone is specific for mono-ADPr and has a preference for Ser-mono-ADPr.				
References	1. Longarini, E.J. <i>et al.</i> (2023) Modular antibodies reveal DNA damage-induced mono-ADP-ribosylation as a second wave of PARP1 signaling. <u>Mol Cell. 83 (10):</u> <u>1743-60.e11.</u>				
Storage	This product is shipped at ambient temperature. It is recommended to aliquot and store at -20°C on receipt. When thawed, aliquot the sample as needed. Keep aliquots at 2-8°C for short term use (up to 4 weeks) and store the remaining aliquots at -20°C.				
	Avoid repeated freezing and thawing as this may denature the antibody. Storage in frost-free freezers is not recommended. Should this product contain a precipitate we recommend microcentrifugation before use.				
Guarantee	12 months from date of despatch				
Acknowledgements	This product and/or its use is covered by claims of U.S. patents, and/or pending U.S. and				

		bio-rad.c	. patent applie com/en-us/trac s a registered	Bio-Rad Laboratories, Inc. See		
	Pailth And SafetyMaterial Safety Datasheet documentation #20479 available aformation https://www.bio-rad-antibodies.com/SDS/TZA020 20479					e at:
Regulato	ry	For rese	arch purpose	s only		
North & South America	Tel: +1 800 265 Fax: +1 919 87 Email: antibody	8 3751	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
To find a b	atch/lot spec	ific datashee	et for this produ	uct, please use our online 'M421387:230706'	search tool at:	bio-rad-antibodies.com/datasheets
				Printed on 28 Jun 2024		
© 2024 Bio-	Rad Laborato	ories Inc L	egal Imprint			