

Datasheet: TZA079SBB580

| | |
|----------------------|---|
| Description: | MOUSE ANTI HUMAN CD66:StarBright Blue 580 |
| Specificity: | CD66 |
| Format: | StarBright Blue 580 |
| Product Type: | Monoclonal Antibody |
| Clone: | rB08-7A10 |
| Isotype: | IgG1 |
| Quantity: | 100 TESTS/0.5ml |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|----------------|-----|----|----------------|--------------------|
| Flow Cytometry | ▪ | | | Neat |

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

| | | | |
|---------------------------------|---|----------------------------|--------------------------|
| Target Species | Human | | |
| Product Form | Purified recombinant IgG containing a SpyTag at the C-terminus of the antibody heavy chain and conjugated to StarBright Blue 580 – liquid | | |
| Max Ex/Em | Fluorophore | Excitation Max (nm) | Emission Max (nm) |
| | StarBright Blue 580 | 475 | 582 |
| Preparation | Purified IgG prepared by affinity chromatography on Protein A from tissue culture supernatant | | |
| Source | CHO cells | | |
| Buffer Solution | Phosphate buffered saline | | |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin 0.1% Pluronic F68 | | |

0.1% PEG 3350

0.05% Tween 20

Approx. Protein Concentrations

For information on the concentration of our StarBright Dye conjugated reagents please visit our [FAQ](#) page.

Immunogen

Recombinant CD66a Gln35-Gly428 expressed by mammalian cells

External Database Links**UniProt:**

[P13688](#) [Related reagents](#)

[P40198](#) [Related reagents](#)

[P06731](#) [Related reagents](#)

[P40199](#) [Related reagents](#)

Entrez Gene:

[634](#) CEACAM1 [Related reagents](#)

[1084](#) CEACAM3 [Related reagents](#)

[1048](#) CEACAM5 [Related reagents](#)

[4680](#) CEACAM6 [Related reagents](#)

Synonyms

BGP, BGP1, CD66D, CEA, CGM1, NCA

Specificity

Mouse anti Human CD66 antibody, clone rB08-7A10, is a recombinant monoclonal antibody that recognizes an epitope common to CD66 a, c, d and e, also known respectively as cell adhesion molecule CEACAM 1, 6, 3 and 5.

CD66 a, c, d and e are members of the carcinoembryonic antigen (CEA) family, which is involved in a range of cellular processes through homotypic and heterotypic interactions. CD66 a, c, d and e all have an extracellular region containing at least one domain homologous to the immunoglobulin variable domain. CD66 a and d are transmembrane proteins, whilst CD66 c and e associate to the cell membrane with GPI anchors. CD66 proteins each have a specific expression pattern. CD66a is broadly expressed and the only family member expressed on activated T cells. CD66a has been attributed to multiple cellular functions, including the modulation of adaptive and innate immune responses, roles in apoptosis, angiogenesis, and involvement in the development of 3D tissue structure. The binding of CD66a by viral and bacterial adhesins can inhibit the function of immune cells, providing a route for pathogens to potentially suppress immune response to infection.

CD66 proteins are associated with immune related diseases and are considered clinical biomarkers and promising therapeutic targets for various types of cancer. CD66a is involved in inflammatory bowel disease, with a protective role in models of colitis. CD66a is a biomarker in non-small cell lung cancer, pancreatic adenocarcinoma and melanoma. CD66e is approved by the FDA as a diagnostic marker for colon cancer.

Flow Cytometry

Use 5µl of the suggested working dilution to label 0.5x10⁶ cells in 100µl. Best practices

suggest a 5 min centrifugation at 6,000g prior to sample application.

| | |
|--------------------------------------|--|
| Storage | Store at +4°C. DO NOT FREEZE. This product should be stored undiluted. This product is photosensitive and should be protected from light. Should this product contain a precipitate we recommend microcentrifugation before use. |
| Guarantee | 12 months from date of despatch |
| Acknowledgements | This product and/or its use is covered by claims of U.S. patents, and/or pending U.S. and non-U.S. patent applications owned by or under license to Bio-Rad Laboratories, Inc. See bio-rad.com/en-us/trademarks for details. This product is covered by U.S Patent No. 10,150,841 and related U.S and foreign counterparts. |
| Health And Safety Information | Material Safety Datasheet documentation #20471 available at: https://www.bio-rad-antibodies.com/SDS/TZA079SBB580 |
| Regulatory | For research purposes only |

Product inquiries: www.bio-rad-antibodies.com/technical-support

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets
'M446665:260109'

Printed on 09 Jan 2026