

## Datasheet: TZC001B

**BATCH NUMBER 168134**

<b>Description:</b>	SpyCatcher2:Biotin
<b>Name:</b>	SPYCATCHER2
<b>Format:</b>	Catcher-Biotin
<b>Product Type:</b>	Recombinant Protein
<b>Quantity:</b>	1 mg

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Immunoassay	▪			

Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Suggested working dilutions are given as a guide only. It is recommended that the user titrates the product for use in their own system using appropriate negative/positive controls.

<b>Target Species</b>	Protein/peptide tag
<b>Product Form</b>	Purified recombinant protein conjugated to Biotin - liquid
<b>Preparation</b>	Recombinant protein expressed in <i>E. coli</i> and purified by affinity chromatography
<b>Source</b>	<i>E. coli</i>
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.0095% MIT
<b>Approx. Protein Concentrations</b>	250 $\mu$ M (3.9 mg/ml)

**Product Information** **SpyCatcher2** (H-SpyC2) is a 15.7 kDa protein that forms a stable covalent isopeptide bond with a second protein that includes a SpyTag. The reaction occurs spontaneously on mixing, is rapid, irreversible, high-yielding and shows good specificity. SpyCatcher2 contains a His6-tag at the N-terminus. To avoid deamidation of asparagine, a N105D

mutation was introduced ([Hentrich et al. 2020](#)).

The reaction is robust at pH 5 to 8, at temperatures from +4°C to +37°C, in various buffer conditions (Ca<sup>2+</sup>/Mg<sup>2+</sup> not needed) and in the presence of detergents. The reaction also occurs inside cells (*in vivo*). Since the speed of the coupling reaction is concentration dependent, it is recommended to use SpyCatcher2 undiluted.

All the different formats of SpyCatcher2 are compatible with Bio-Rad's recombinant HuCAL® antibodies with a SpyTag2 at the C-terminus of the heavy chain.

[Download SpyCatcher coupling protocol](#)

SpyCatcher2 (TZC001) is available conjugated to biotin (TZC001B). It is also available as a modified version, SpyCatcher2-CYS (TZC001CYS), which contains a cysteine residue for site-specific conjugation by the user.

SpyCatcher2-CYS can dimerize during storage by formation of a disulfide bond via the free cysteines. Before conjugation to these cysteines, the SpyCatcher2-CYS dimers should be mildly reduced e.g., by addition of 5mM DTT and incubation for 1 hr at room temperature, followed by a fast DTT removal step, e.g., by size exclusion chromatography using PD-10 desalting columns, as DTT can interfere with the subsequent conjugation chemistry. Dialysis is not recommended for DTT removal, as the cysteines can oxidize again during this slow process.

[View all available Catchers](#)

---

<b>Protein Molecular Weight</b>	Predicted molecular weight of the purified Catcher is 15904 Da The molecular weight specified excludes conjugation and will vary upon conjugation depending on the molecular weight of the label
<b>Purity</b>	≥90% determined by SDS-PAGE under reducing conditions and visualized by coomassie blue staining
<b>Instructions For Use</b>	<a href="#">View SpyCatcher Coupling Protocol</a>
<b>References</b>	1. Keeble, A.H. <i>et al.</i> (2017) Evolving Accelerated Amidation by SpyTag/SpyCatcher to Analyze Membrane Dynamics. <a href="#">Angew Chem Int Ed Engl. 56 (52): 16521-5.</a> 2. Hentrich, C. <i>et al.</i> (2021) Periplasmic expression of SpyTagged antibody fragments enables rapid modular antibody assembly. <a href="#">Cell Chem Biol. 28 (6): 813-824.e6.</a>
<b>Storage</b>	Store at -20°C only. Storage in frost-free freezers is not recommended. This product should be stored undiluted. Avoid repeated freezing and thawing as this may denature the protein. Should this product contain a precipitate we recommend microcentrifugation before use.
<b>Guarantee</b>	12 months from date of despatch

---

**Acknowledgements** This product and/or its use is covered by claims of U.S. patents, and/or pending U.S. and non-U.S. patent applications owned by or under license to Bio-Rad Laboratories, Inc. See [bio-rad.com/en-us/trademarks](http://bio-rad.com/en-us/trademarks) for details.  
His-tag is a registered trademark of EMD Biosciences.

---

**Health And Safety Information** Material Safety Datasheet documentation #20480 available at: <https://www.bio-rad-antibodies.com/SDS/TZC001B>  
20480

---

**Licensed Use** For *in vitro* research purposes. Any re-sale in any form or any other commercial application needs a written agreement with Bio-Rad.

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Useful Reagents

[SpyCatcher2 \(TZC001\)](#)

[SpyCatcher2-CYS \(TZC001CYS\)](#)

[ANTI SpyCatcher \(HCA379\)](#)

[SpyTag3 PEPTIDE \(BLP086\)](#)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](http://bio-rad-antibodies.com/datasheets)  
'M422625:230921'

**Printed on 17 Apr 2024**

---

© 2024 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)