

## Datasheet: VMA00016KT

<b>Description:</b>	POLY (ADP-RIBOSE) POLYMERASE 1 ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	POLY (ADP-RIBOSE) POLYMERASE 1
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb™ Monoclonal
<b>Isotype:</b>	IgG1
<b>Quantity:</b>	2 Westerns

## Product Details

### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Species Cross Reactivity</b>	Reacts with: Mouse, Rat <b>N.B.</b> Antibody reactivity and working conditions may vary between species.
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Immunogen</b>	Human PARP-1
<b>External Database Links</b>	<p><b>UniProt:</b>  <a href="#">P09874</a>   <a href="#">Related reagents</a></p> <p><b>Entrez Gene:</b>  <a href="#">142</a> PARP1   <a href="#">Related reagents</a></p>
<b>Synonyms</b>	ADPRT, PPOL

<b>Specificity</b>	<p><b>Mouse anti Human poly (ADP-ribose) polymerase 1 antibody</b> recognizes poly (ADP-ribose) polymerase 1 (PARP-1), a nuclear enzyme cleaved during apoptosis (<a href="#">Soldani et al. 2002</a>).</p> <p>PARP-1 is a caretaker enzyme involved in DNA damage repair (<a href="#">Langelier et al. 2013</a>). It plays roles in diabetes pathophysiology (<a href="#">Andreone et al. 2012</a>) and tumor proliferation (<a href="#">Rosado et al. 2013</a>). As well as protecting cells from genomic instability, PARP-1 is also involved in the development of both inflammatory and immune responses, and cell death by apoptosis and necrosis (<a href="#">Erdélyi et al. 2005</a>).</p> <p>PARP-1 represents a promising target for new developments in therapeutic treatment of immune mediated diseases (<a href="#">Rosado et al. 2013</a>) and has considerable potential for delivering selective tumor cell killing while sparing normal cells (<a href="#">Pinton et al. 2013</a>).</p> <p>Mouse anti Human poly (ADP-ribose) polymerase 1 antibody recognizes PARP-1 as a single band of ~116 kDa in western blotting in multiple cell line lysates including myeloid, lymphoid, neuronal, adenocarcinoma, sarcoma and embryonic kidney.</p>
<b>Western Blotting</b>	Anti Poly (ADP-Ribose) Polymerase 1 antibody detects a band of approximately 116 kDa in HEK293 cell lysates.
<b>Instructions For Use</b>	Please refer to the <a href="#">PrecisionAb western blotting protocol</a> . For additional information on secondary antibody dilution and exposure time see product web page.
<b>Lysate Composition</b>	400µg HEK293 lysate lyophilized in RIPA buffer.
<b>Lysate Reconstitution</b>	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
<b>Storage</b>	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
<b>Guarantee</b>	As supplied, 12 months from date of despatch.
<b>Acknowledgements</b>	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
<b>Health And Safety Information</b>	<p>Material Safety Datasheet documentation #10040 #10561 available at:</p> <p>Antibody (10040): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</a></p> <p>Lysate Material (10561): <a href="https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf">https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</a></p>
<b>Regulatory</b>	For research purposes only

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

## Recommended Negative Controls

### MOUSE IgG1 NEGATIVE CONTROL (MCA928)

**North & South** Tel: +1 800 265 7376

**America** Fax: +1 919 878 3751

Email: [antibody\\_sales\\_us@bio-rad.com](mailto:antibody_sales_us@bio-rad.com)

**Worldwide**

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: [antibody\\_sales\\_uk@bio-rad.com](mailto:antibody_sales_uk@bio-rad.com)

**Europe**

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: [antibody\\_sales\\_de@bio-rad.com](mailto:antibody_sales_de@bio-rad.com)

'M338052:181217'

**Printed on 15 Mar 2019**

---

© 2019 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)