

Datasheet: VMA00021KT

Description:	CD147 ANTIBODY WITH CONTROL LYSATE
Specificity:	CD147
Other names:	Neurothelin
Format:	Purified
Product Type:	PrecisionAb™ Monoclonal
Isotype:	lgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Human		
Purified IgG - liquid		
20μl Mouse monoclonal antibody purified by affinity chromatography on Protein A from ascites		
Phosphate buffered saline		
0.09% Sodium Azide (NaN ₃)		
Soluble recombinant form of CD147 (CD147Rg) consisting of the cDNA coding for the hinge region, CH2 and CH3 domain of human IgG1		
UniProt: P35613 Related reagents Entrez Gene: 682 BSG Related reagents		

Specificity

Mouse anti Human CD147 antibody recognizes the human CD147 cell surface antigen, also known as Basigin, EMMPRIN and collagenase stimulatory factor. CD147 is a 385 amino acid single

pass type 1 trans-membrane glycoprotein bearing 3 potential N-glycosylation sites, a single N-terminal (distal) <u>Ig-like C2-type domain</u> and a more proximal <u>Ig-like V-type domain</u> in its extracellular region. Mouse anti Human CD147 antibody was raised and screened against recombinant extracellular region of huCD147 expressed in CHO cells. Epitope mapping indicates that this antibody binds to an epitope within the distal <u>Ig-like C-2</u> type domain of human CD147.

Mouse anti Human CD147 antibody binds to both un-stimulated and phytohemagglutinin (PHA) stimulated (activated) T lymphocytes with levels of CD147 enhanced following PHA stimulation. CD147 is also expressed by peripheral blood and endothelial cells as well as many cultured cells of hematopoietic and non-hematopoietic origin.

Mouse anti Human CD147 antibody recognizes human CD147 as a band of ~53 kDa by western blot analysis of HepG2 cell lysates under reducing conditions.

Western Blotting	Anti CD147 antibody detects a band of approximately 53 kDa in HepG2 cell lysates.
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg HepG2 lysate lyophilized in RIPA buffer.
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 10 μ l 2M DTT.
	- If using BME reconstitute the lyophilized lysate with 180 μ l DI H $_2$ O, add 200 μ l 2x Laemmli Sample Buffer and 20 μ l BME.
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Shelf Life	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf

Related Products

Regulatory

Recommended Secondary Antibodies

For research purposes only

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

'M336496:181214'

North & South Tel: +1 800 265 7376 America Fax: +1 919 878 3751 Worldwide

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_uk@bio-rad.com Email: antibody_sales_de@bio-rad.com

Printed on 02 Jan 2019

© 2019 Bio-Rad Laboratories Inc | Legal | Imprint

Email: antibody_sales_us@bio-rad.com