

Datasheet: VMA00043 BATCH NUMBER 080714

Description: MOUSE ANTI VISFAT		
Specificity:	VISFATIN	
Other names:	NAMPT	
Format:	Purified	
Product Type:	PrecisionAb Monoclonal	
Isotype:	lgG1	
Quantity:	100 μΙ	

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody purified by affinity chromatography on Protein G from ascites
Buffer Solution	Phosphate buffered saline
Preservative	0.09% Sodium Azide (NaN ₃)

Stabilisers

cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2006). It has also been suggested that macrophages in visceral white adipositissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1α and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under						
Links UniProt: P43490 Related reagents Entrez Gene: 10135 NAMPT Related reagents Synonyms PBEF, PBEF1 Specificity Mouse anti Human visfatin antibody recognizes human visfatin also known as nicotinamide phosphoribosyltransferase and pre-B-cell colony-enhancing factor 1, a putative cytokine and member of the NAPRTase family involved in catalyzing the biosynthesis of NAD from nicotinamide. Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendritic cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2006). It has also been suggested that macrophages in visceral white adiportissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1c and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at ~20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)	Immunogen	Recombinant human visfatin, amino acids 1-491, purified from <i>E. coli</i>				
Entrez Gene: 10135 NAMPT Related reagents Synonyms PBEF, PBEF1 Specificity Mouse anti Human visfatin antibody recognizes human visfatin also known as nicotinamide phosphoribosyltransferase and pre-B-cell colony-enhancing factor 1, a putative cytokine and member of the NAPRTase family involved in catalyzing the biosynthesis of NAD from nicotinamide. Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendriticells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2006). It has also been suggested that macrophages in visceral white adipotissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1a and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at ~20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch PrecisionAb is a trademark of Bio-Rad Laboratories. Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)		UniProt:				
Synonyms PBEF, PBEF1 Specificity Mouse anti Human visfatin antibody recognizes human visfatin also known as nicotinamide phosphoribosyltransferase and pre-B-cell colony-enhancing factor 1, a putative cytokine and member of the NAPRTase family involved in catalyzing the biosynthesis of NAD from nicotinamide. Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendritic cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2008). It has also been suggested that macrophages in visceral white adiportissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1α and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)		P43490 Related reagents				
Synonyms PBEF, PBEF1 Specificity Mouse anti Human visfatin antibody recognizes human visfatin also known as nicotinamide phosphoribosyltransferase and pre-B-cell colony-enhancing factor 1, a putative cytokine and member of the NAPRTase family involved in catalyzing the biosynthesis of NAD from nicotinamide. Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendritic cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2006). It has also been suggested that macrophages in visceral white adiport tissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1α and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at ~20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch PrecisionAb is a trademark of Bio-Rad Laboratories. Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)		Entrez Gene:				
Mouse anti Human visfatin antibody recognizes human visfatin also known as nicotinamide phosphoribosyltransferase and pre-B-cell colony-enhancing factor 1, a putative cytokine and member of the NAPRTase family involved in catalyzing the biosynthesis of NAD from nicotinamide. Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendritic cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabeted realities (Chen et al. 2006). It has also been suggested that macrophages in visceral white adipositissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1a and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)		10135 NAMPT Related reagents				
nicotinamide phosphoribosyltransferase and pre-B-cell colony-enhancing factor 1, a putative cytokine and member of the NAPRTase family involved in catalyzing the biosynthesis of NAD from nicotinamide. Visfatin is highly expressed in bone marrow, muscle and liver tissue and increases in expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendritic cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Eukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2006). It has also been suggested that macrophages in visceral white adiportissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1α and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch PrecisionAb is a trademark of Bio-Rad Laboratories. Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)	Synonyms	PBEF, PBEF1				
expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendritic cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produce by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2006). It has also been suggested that macrophages in visceral white adiportissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1α and IL-8, suggesting a pro-inflammatory marker role for visfatin (Curat et al. 2006). Mouse anti Human Visfatin antibody recognizes human visfatin as a single band of ~56 kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch PrecisionAb is a trademark of Bio-Rad Laboratories. Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)	Specificity	nicotinamide phosphoribosyltransferase and pre-B-cell colony-enhancing factor 1, a putative cytokine and member of the NAPRTase family involved in catalyzing the				
kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell lines. Western Blotting Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates. Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)		expression in response to endotoxin, in amniotic epithelial cells following lipopolysaccharide (LPS) stimulation, and during the activation of T- and B-cells, dendritic cells and macrophages, suggesting a role in inflammatory processes. Visfatin is produced by visceral fat and appears to have effects similar to those of insulin (Fukuhara et al. 2005). Plasma levels of visfatin are elevated in patients with type 2 diabetes mellitus (Chen et al. 2006). It has also been suggested that macrophages in visceral white adipose tissue may be a prominent source of the adipokines visfatin and resistin as well as chemokines MCP-1, MIP1α and IL-8, suggesting a pro-inflammatory marker role for				
Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)		kDa in K562 cell lysates and in other human cell line lysates by western blotting under reducing conditions. The antibody also cross-reacts with visfatin from mouse and rat cell				
Guarantee 12 months from date of despatch Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)	Western Blotting	Anti visfatin antibody detects a band of approximately 56 kDa in K562 cell lysates.				
Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories. Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)	Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.				
Health And Safety Information Material Safety Datasheet documentation #10040 available at: https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)	Guarantee	12 months from date of despatch				
Information https://www.bio-rad-antibodies.com/SDS/VMA00043 Antibody (10040)	Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.				
Regulatory For research purposes only		https://www.bio-rad-antibodies.com/SDS/VMA00043				
	Regulatory	For research purposes only				

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

 $\label{lem:lemail:antibody_sales_uk@bio-rad.com} \\ Email: antibody_sales_uk@bio-rad.com \\ Email: antibody_sales_uk@bio-rad.com \\ \\ Email: antibody_sales_de@bio-rad.com \\ \\ \\ Email: antibody_s$

To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M369862:200529'

Printed on 13 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint