

Datasheet: VMA00070KT

Description:	CA125 ANTIBODY WITH CONTROL LYSATE
Specificity:	CA125
Other names:	MUCIN 16
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	lgG1
Quantity:	2 Westerns

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .							
		Yes	No	Not Determined	Suggested Dilution			
	Western Blotting	-			1/1000			
	application. The product has not	e antibody has bee been tested for us	n validate e in a par	ively <u>validated for th</u>ed at the suggested di ticular technique this o optimization may be re	lution. Where this			
Target Species	Human							
Product Form	Purified IgG - liq	uid						
Preparation	20µl Mouse mor	noclonal antibody p	urified by	ion exchange chroma	atography from ascites			
Buffer Solution	Phosphate buffered saline							
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)							
Immunogen	CA125 antigen purified from human ovarian carcinoma							
External Database Links	UniProt: <u>Q8WXI7</u>	Related reagents						

Entrez Gene:

94025 MUC16 Related reagents

Synonyms	CA125
Specificity	Mouse anti Human CA125 antibody recognizes the CA125, also known as mucin-16, ovarian cancer-related tumor marker CA125 or ovarian carcinoma antigen CA125.
	Mouse anti Human CA125 antibody detects a band of 2253 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti CA125 detects a band of approximately 2253 kDa in HeLa cell lysates.
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg HeLa lysate lyophilized in RIPA buffer.
Lysate Reconstitution	 If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21	
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50	From
	Email: antibody_sales_us@bio-rad.com		Email: antibody_sales_uk@bio-rad.com		Email: antibody_sales_de@bio-rad.mmrch	

15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M350937:190315'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint