

Datasheet: VMA00094KT

Description:	PRO DESMOGLEIN 2 ANTIBODY WITH CONTROL LYSATE
Specificity:	PRO DESMOGLEIN 2
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	20G1
Isotype:	IgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000
Immunofluorescence		▪		

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein A from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Recombinant fusion protein fragment consisting of amino acids 37-164 of desmoglein 2 which consists of the entire extracellular domain and a portion of the proregion.

**External Database
Links**

UniProt:

[Q14126](#) [Related reagents](#)

Entrez Gene:

[1829](#) DSG2 [Related reagents](#)

Synonyms

CDHF5

Specificity

Mouse anti Human pro desmoglein 2 recognizes pro desmoglein 2, a member of the desmoglein family. Desmosomes are cell-cell junctions between epithelial, myocardial, and certain other cell types. Currently, four desmoglein subfamily members have been identified and all are members of the cadherin cell adhesion molecule superfamily.

Desmoglein 2 is a membrane glycoprotein with 4 highly conserved extracellular domains (EC1 - 4) and six cytoplasmic desmoglein repeat sequences. It has a predicted molecular weight of 122 kDa and an apparent molecular weight of 150 kDa, due to post translational modification. Desmoglein 2 is a core component of the desmosome cell-cell junction between cells. This protein has been identified as the autoantigen of the autoimmune skin blistering disease pemphigus vulgaris and mutations in the gene have been associated with arrhythmogenic right ventricular dysplasia.

Mouse anti Human pro desmoglein 2 specifically recognizes an epitope within the proregion domain of desmoglein 2, epitope mapping having demonstrated reactivity with fusion proteins containing the entire prodomain but not with fusion proteins lacking the proregion ([Keim et al., 2008](#)). This antibody has been extensively validated for western blotting using whole cell lysates and is available with a positive control lysate, in a smaller trial size, and as a standalone antibody.

Western Blotting

Anti pro desmogein 2 antibody detects a band of approximately 150 kDa in HepG2 cell lysates

Instructions For Use

Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

Lysate Composition

400µg HepG2 lysate lyophilized in RIPA buffer.

Lysate Reconstitution

- If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.
- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.
Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.

References

1. Keim, S.A. *et al.* (2008) Generation and characterization of monoclonal antibodies against the proregion of human desmoglein-2. [Hybridoma 27: 249-58.](#)

Storage

Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.

Guarantee	As supplied, 12 months from date of despatch.
------------------	---

Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
-------------------------	--

Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
--------------------------------------	--

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at bio-rad-antibodies.com

'M351458:190318'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)