

Datasheet: VMA00105

BATCH NUMBER 290715

Description:	MOUSE ANTI CYSTATIN C
Specificity:	CYSTATIN C
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	IgG1
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Yeast-derived recombinant protein
External Database Links	UniProt: P01034 Related reagents

Entrez Gene:[1471](#) CST3 [Related reagents](#)**Specificity**

Mouse anti Human cystatin C antibody recognizes cystatin C, also known as bA218C14.4, cystatin 3, cystatin-3, gamma-trace, neuroendocrine basic polypeptide and post-gamma-globulin.

The cystatin superfamily encompasses proteins that contain multiple cystatin-like sequences. Some of the members are active cysteine protease inhibitors, while others have lost or perhaps never acquired this inhibitory activity. There are three inhibitory families in the superfamily, including the type 1 cystatins (stefins), type 2 cystatins and the kininogens. The cystatin locus on chromosome 20 contains the majority of the type 2 cystatin genes and pseudogenes. The CST3 gene, which encodes Cystatin C, is located in the cystatin locus and encodes the most abundant extracellular inhibitor of cysteine proteases, which is found in high concentrations in biological fluids and is expressed in virtually all organs of the body. A mutation in CST3 has been associated with amyloid angiopathy. Expression of this protein in vascular wall smooth muscle cells is severely reduced in both atherosclerotic and aneurysmal aortic lesions, establishing its role in vascular disease (provided by RefSeq, Jul 2008).

Mouse anti Human cystatin C antibody detects a band of 13 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting

Anti cystatin C detects a band of approximately 13 kDa in A431 cell lysates

Storage

Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Guarantee

12 months from date of despatch

Acknowledgements

PrecisionAb is a trademark of Bio-Rad Laboratories.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:
<https://www.bio-rad-antibodies.com/SDS/VMA00105>
Antibody (10040)

Regulatory

For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

North & South America

Tel: +1 800 265 7376

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

To find a batch/lot specific datasheet for this product, please use our online search tool at: [bio-rad-antibodies.com/datasheets](https://www.bio-rad-antibodies.com/datasheets)

'M369906:200529'

