

Datasheet: VMA00124

Description:	MOUSE ANTI RAF-1
Specificity:	RAF-1
Other names:	Proto-oncogene c-RAF
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	IgG1
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Purified IgG prepared by affinity chromatography on Protein G from ascites
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Purified recombinant fragment of human RAF-1 expressed in <i>E. coli</i>
External Database Links	<p>UniProt: P04049 Related reagents</p> <p>Entrez Gene: 5894 RAF1 Related reagents</p>

Synonyms RAF

Specificity **Mouse anti Human RAF-1 antibody** recognizes RAF-1, also known as C-Raf proto-oncogene, serine/threonine kinase, Oncogene RAF1, RAF proto-oncogene serine/threonine-protein kinase, proto-oncogene c-RAF, raf proto-oncogene serine/threonine protein kinase or v-raf-1 murine leukemia viral oncogene homolog 1.

RAF-1 is the cellular homolog of viral raf gene (v-raf). The encoded protein is a MAP kinase kinase kinase (MAP3K), which functions downstream of the Ras family of membrane associated GTPases to which it binds directly. Once activated, the cellular RAF-1 protein can phosphorylate to activate the dual specificity protein kinases MEK1 and MEK2, which in turn phosphorylate to activate the serine/threonine specific protein kinases, ERK1 and ERK2. Activated ERKs are pleiotropic effectors of cell physiology and play an important role in the control of gene expression involved in the cell division cycle, apoptosis, cell differentiation and cell migration. Mutations in RAF-1 are associated with Noonan syndrome 5 and LEOPARD syndrome 2 (provided by RefSeq, Jul 2008).

Mouse anti Human RAF-1 antibody detects a band of 73 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting Anti RAF-1 detects a band of approximately 73 kDa in HEK293 cell lysates.

Further Reading 1. Morrison, D.K. & Cutler, R.E. (1997) The complexity of Raf-1 regulation. [Curr Opin Cell Biol. 9\(2\):174-9](#)

Storage Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Guarantee 12 months from date of despatch

Acknowledgements PrecisionAb is a trademark of Bio-Rad Laboratories.

Health And Safety Information Material Safety Datasheet documentation #10040 available at: Antibody (10040): <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M369920:200529'

Printed on 27 Sep 2020