

Datasheet: VMA00153

Description:	MOUSE ANTI GLUTAMATE DECARBOXYLASE 2
Specificity:	GLUTAMATE DECARBOXYLASE 2
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	IgG1
Quantity:	100 µl

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	■			1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click [here](#) to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species.
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)
Immunogen	Keyhole limpet haemocyanin (KLH) conjugated synthetic peptide sequence PGSGFWSFGSEdGSGDSEN corresponding to amino acids 4-22 within the N-terminal region of human glutamate decarboxylase 2
External Database Links	UniProt: Q05329 Related reagents Entrez Gene:

Synonyms	GAD65
-----------------	-------

Specificity	<p>Mouse anti Human glutamate decarboxylase 2 antibody recognizes glutamate decarboxylase 2, also known as 65 kDa glutamic acid decarboxylase, GAD-65, or glutamate decarboxylase 65 kDa isoform.</p> <p>Encoded by the GAD2 gene, glutamate decarboxylase 2 is one of several forms of glutamic acid decarboxylase. Glutamate decarboxylase 2 is responsible for catalyzing the production of gamma-aminobutyric acid from L-glutamic acid. A pathogenic role for this enzyme has been identified in the human pancreas since it has been identified as an autoantibody and an autoreactive T cell target in insulin-dependent diabetes. GAD2 may also play a role in the stiff man syndrome. Alternative splicing results in multiple transcript variants that encode the same protein (provided by RefSeq, Oct 2008).</p> <p>Mouse anti Human glutamate decarboxylase 2 antibody detects a band of 60 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
--------------------	---

Western Blotting	Anti glutamate Decarboxylase 2 detects a band of approximately 60 kDa in Mouse Brain cell lysates.
-------------------------	--

Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
----------------	---

Guarantee	12 months from date of despatch
------------------	---------------------------------

Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.
-------------------------	---

Health And Safety Information	Material Safety Datasheet documentation #10040 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf
--------------------------------------	---

Regulatory	For research purposes only
-------------------	----------------------------

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South Tel: +1 800 265 7376

America Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Worldwide

Tel: +44 (0)1865 852 700

Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21

Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

'M369946:200529'

Printed on 11 Aug 2020