

Datasheet: VMA00167KT

| Description: | FOCAL ADHESION KINASE ANTIBODY WITH CONTROL LYSATE |
|---------------|--|
| Specificity: | FOCAL ADHESION KINASE |
| Format: | Purified |
| Product Type: | PrecisionAb Monoclonal |
| Clone: | OTI4A8 |
| Isotype: | lgG1 |
| Quantity: | 2 Westerns |
| | |

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

| | Yes | No | Not Determined | Suggested Dilution |
|------------------|-----|----|----------------|--------------------|
| Western Blotting | | | | 1/1000 |

PrecisionAb antibodies have been extensively <u>validated for the western blot</u> <u>application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

| Target Species | Human |
|-----------------------------|--|
| Species Cross Reactivity | Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. |
| Product Form | Purified IgG - liquid |
| Preparation | 20μl Mouse monoclonal antibody purified by affinity chromatography from ascites. |
| Buffer Solution | Phosphate buffered saline |
| Preservative Stabilisers | 0.09% Sodium Azide (NaN ₃) 1% Bovine Serum Albumin |

| | 50% Glycerol | | | |
|----------------------------|---|--|--|--|
| Immunogen | Full length recombinant protein of human focal adhesion kinase (NP_722560) produced in HEK293T cells | | | |
| External Database Links | UniProt: Q05397 Related reagents Entrez Gene: 5747 PTK2 Related reagents | | | |
| Synonyms | FAK, FAK1 | | | |
| Specificity | Mouse anti Human focal adhesion kinase 1 antibody recognizes focal adhesion kinase 1, also known as FADK 1, FAK-related non-kinase polypeptide, PTK2 protein tyrosine kinase 2, focal adhesion kinase 1, focal adhesion kinase isoform FAK-Del33, focal adhesion kinase-related nonkinase, protein phosphatase 1 regulatory subunit 71 and protein phosphatase 1, regulatory subunit 71. Encoded by the PTK2 gene, focal adhesion kinase 1 is a cytoplasmic protein tyrosine kinase which is found concentrated in the focal adhesions that form between cells growing in the presence of extracellular matrix constituents. The encoded protein is a member of the FAK subfamily of protein tyrosine kinases but lacks significant sequence similarity to kinases from other subfamilies. Activation of focal adhesion kinase 1 may be an important early step in cell growth and intracellular signal transduction pathways triggered in response to certain neural peptides or to cell interactions with the extracellular matrix. Several transcript variants encoding different isoforms have been found for PTK2, but the full-length natures of only three of them have been determined (provided by RefSeq, Dec 2010). Mouse anti Human focal adhesion kinase 1 antibody detects a band of 125 kDa. The | | | |
| Western Blotting | antibody has been extensively validated for western blotting using whole cell lysates. Anti-focal adhesion kinase detects a hand of approximately 125 kDa in A549 cell lysates. | | | |
| Instructions For Use | Anti focal adhesion kinase detects a band of approximately 125 kDa in A549 cell lysates. Please refer to the PrecisionAb western blotting protocol. For additional information on | | | |
| instructions for use | secondary antibody dilution and exposure time see product web page. | | | |
| Lysate Composition | 400μg A549 lysate lyophilized in RIPA buffer. | | | |
| Lysate Reconstitution | - If using DDT reconstitute the lyophilized lysate with 190 μ l DI H ₂ O, add 200 μ l 2x Laemmli Sample Buffer and 10 μ l 2M DTT If using BME reconstitute the lyophilized lysate with 180 μ l DI H ₂ O, add 200 μ l 2x Laemmli Sample Buffer and 20 μ l BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25 μ l. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. | | | |

| Storage | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. | | | | |
|----------------------------------|--|--|--|--|--|
| | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. | | | | |
| Guarantee | As supplied, 12 months from date of despatch. | | | | |
| Acknowledgements | PrecisionAb™ is a trademark of Bio-Rad Laboratories. | | | | |
| Health And Safety Information | Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf | | | | |
| Regulatory | For research purposes only | | | | |

Related Products

America

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376 Worldwide

Fax: +1 919 878 3751

Email: antibody_sales_us@bio-rad.com

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739

Email: antibody_sales_uk@bio-rad.com

Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351136:190315'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint