

# Datasheet: VMA00176KT

Description:	BTK ANTIBODY WITH CONTROL LYSATE				
Specificity:	ВТК				
Format:	Purified				
Product Type:	PrecisionAb Monoclonal				
Clone:	OTI12D4				
lsotype:	lgG2a				
Quantity:	2 Westerns				

# **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .						
		Yes	No	Not Determined	Suggested Dilution		
	Western Blotting	•			1/1000		
	PrecisionAb antibodies application. The antibod product has not been tes exclude its use in such p sample type.	dy has bee sted for use	n validate e in a par	ed at the suggested di ticular technique this c	lution. Where this does not necessarily		
Target Species	Human						
Product Form	Purified IgG - liquid						
Preparation	20µl Mouse monoclonal antibody purified by affinity chromatography from ascites.						
Buffer Solution	Phosphate buffered salir	ie					
Preservative	0.09% Sodium Azide (Na	aN <sub>3</sub> )					
Stabilisers	1% Bovine Serum Album	nin					
	50% Glycerol						
Immunogen	Full length recombinant l	human BT	K (NP_00	00052) produced in HE	EK293T cells		
External Database Links	UniProt:						

	Q06187 Related reagents						
	Entrez Gene: 695 BTK <u>Related reagents</u>						
Synonyms	AGMX1, ATK, BPK						
Specificity	<b>Mouse anti Human BTK antibody</b> recognizes BTK, also known as B-cell progenitor kinase, gamma globulinaemia tyrosine kinase, dominant-negative kinase-deficient Brutons tyrosine kinase, truncated Bruton gamma globulinemia tyrosine kinase, tyrosine-protein kinase BTK and tyrosine-protein kinase BTK isoform (lacking exon 14).						
	The protein encoded by the BTK gene plays a crucial role in B-cell development. Mutations in BTK cause X-linked agammaglobulinemia type 1, which is an immunodeficiency characterized by the failure to produce mature B lymphocytes, and associated with a failure of Ig heavy chain rearrangement (provided by RefSeq, Nov 2008).						
	Mouse anti Human BTK antibody detects a band of 72 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.						
Western Blotting	Anti BTK detects a band of approximately 72 kDa in K562 cell lysates.						
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.						
Lysate Composition	400µg K562 lysate lyophilized in RIPA buffer.						
Lysate Reconstitution	- If using DDT reconstitute the lyophilized lysate with 190µl DI H <sub>2</sub> O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.						
	- If using BME reconstitute the lyophilized lysate with 180µl DI H <sub>2</sub> O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.						
	Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.						
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.						
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.						
Guarantee	As supplied, 12 months from date of despatch.						
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.						
Health And Safety Information	Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>						

## **Related Products**

### **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (H/L) (STAR207...)<u>HRP</u> Recommended Negative Controls

#### MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide	Tel: +44 (0)1865 852 700	Europe	Tel: +49 (0) 89 8090 95 21
America	Fax: +1 919 878 3751		Fax: +44 (0)1865 852 739		Fax: +49 (0) 89 8090 95 50
	Email: antibody_sales_us@bio-ra	id.com	Email: antibody_sales_uk@bio-ra	ad.com	Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M350940:190315'

### Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint