

Datasheet: VMA00201KT

Description:	CD146 ANTIBODY WITH CONTROL LYSATE
Specificity:	CD146
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	OTI2G10
Isotype:	IgG2b
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting				1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot</u> <u>application.</u> The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

External Database Links	UniProt:
Immunogen	Full length recombinant human CD146 (NP_006491) produced in HEK293T cells
	50% Glycerol
Stabilisers	1% Bovine Serum Albumin
Preservative	0.09% Sodium Azide (NaN ₃)
Buffer Solution	Phosphate buffered saline
Preparation	20μl Mouse monoclonal antibody purified by affinity chromatography from ascites.
Product Form	Purified IgG - liquid
Target Species	Human

P43121	Related reagents

Entrez Gene:

4162 MCAM Related reagents

Synonyms	MUC18
Specificity	Mouse anti Human CD146 antibody recognizes CD146, also known as gicerin, S-endo 1 endothelial-associated antigen, cell surface glycoprotein MUC18, cell surface glycoprotein P1H12, melanoma adhesion molecule, melanoma-associated antigen A32 and melanoma-associated antigen MUC18.
	Mouse anti Human CD146 antibody detects a band of 120 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti CD146 detects a band of approximately 120 kDa in HeLa cell lysates.
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol</u> . For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg HeLa lysate lyophilized in RIPA buffer.
Lysate Reconstitution	 If using DDT reconstitute the lyophilized lysate with 190μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT. If using BME reconstitute the lyophilized lysate with 180μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 20μl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10048 #10561 available at: Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG2b NEGATIVE CONTROL (MCA691)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351770:190318'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint