

## Datasheet: VMA00234

Description:	MOUSE ANTI OCT4
Specificity:	OCT4
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	OTI2B6
lsotype:	lgG2b
Quantity:	100 µl

## **Product Details**

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .					
		Yes	No	Not Determined	Suggested Dilution	
	Western Blotting				1/1000	
	The PrecisionAb label is reserved for antibodies that meet the defined performance					
	criteria within Bio-Rad's how we validate our Prec use in a particular techniqu Further optimization may b	<b>cisionAb</b> ue this do	<b>range.</b> V es not no	Where this product has ecessarily exclude its	s not been tested for	
Target Species	Human					
Product Form	Purified IgG - liquid					
Preparation	Mouse monoclonal antiboo	dy purified	l by affin	ity chromatography fro	om ascites.	
Buffer Solution	Phosphate buffered saline.	•				
Preservative	0.09% Sodium Azide (NaN	l <sub>3</sub> )				
Stabilisers	1% Bovine Serum Albumin					
	50% Glycerol.					
Immunogen	Full length recombinant hu	iman OCT	Γ4 (NP_(	002692) produced in H	HEK293T cells	
External Database Links	UniProt: <u>Q01860</u> <u>Related rea</u>	agents				

	Entrez Gene: 5460 POU5F1 Related reagents				
Synonyms	OCT3, OCT4, OTF3				
Specificity	<b>Mouse anti Human OCT4 antibody, clone OTI2B6</b> recognizes the POU domain, class 5, transcription factor 1, also known as POU domain transcription factor OCT4, POU domain, class 5, transcription factor 1, POU-type homeodomain-containing DNA-binding protein, octamer-binding protein 3, octamer-binding protein 4 and octamer-binding transcription factor 3.				
	The OCT4 gene encodes a transcription factor containing a POU homeodomain. This transcription factor plays a role in embryonic development, especially during early embryogenesis, and it is necessary for embryonic stem cell pluripotency. A translocation of POU5F1 with the Ewing's sarcoma gene, t(6;22)(p21;q12), has been linked to tumor formation. Alternative splicing, as well as usage of alternative translation initiation codons, results in multiple isoforms, one of which initiates at a non-AUG (CUG) start codon. Related pseudogenes have been identified on chromosomes 1, 3, 8, 10, and 12 (provided by RefSeq, Mar 2010).				
	antibody has been extensively validated for western blotting using whole cell lysates.				
Western Blotting	Anti OCT4 detects a band of approximately 45 kDa in NTERA-2 cell lysates.				
Storage	Store undiluted at -20°C, avoiding repeated freeze thaw cycles.				
Guarantee	12 months from date of despatch.				
Acknowledgements	PrecisionAb is a trademark of Bio-Rad Laboratories.				
Health And Safety Information	Material Safety Datasheet documentation #10048 available at: https://www.bio-rad-antibodies.com/SDS/VMA00234 Antibody (10048)				

## **Recommended Secondary Antibodies**

Goat Anti Mouse IgG (H/L) (STAR207...) <u>HRP</u> Recommended Negative Controls

MOUSE IgG2b NEGATIVE CONTROL (MCA691)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad	.com

Tel: +44 (0)1865 852 700 **Europe** Fax: +44 (0)1865 852 739 Email: antibody\_sales\_uk@bio-rad.com Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody\_sales\_de@bio-rad.com To find a batch/lot specific datasheet for this product, please use our online search tool at: bio-rad-antibodies.com/datasheets 'M416470:230301'

## Printed on 13 Aug 2023

© 2023 Bio-Rad Laboratories Inc | Legal | Imprint