

Datasheet: VMA00289KT

Description:	HISTONE DEACETYLASE 4 ANTIBODY WITH CONTROL LYSATE
Specificity:	HISTONE DEACETYLASE 4
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	lgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	•			1/1000

PrecisionAb antibodies have been extensively <u>validated for the western blot</u> <u>application</u>. The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human		
Product Form	Purified IgG - liquid		
Preparation	20μl Mouse monoclonal antibody prepared by affinity chromatography on Protein G from ascites		
Buffer Solution	Phosphate buffered saline		
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)		
Immunogen	Purified recombinant fragment of human histone deacetylase 4 expressed in E. coli		
External Database Links	UniProt: P56524 Related reagents		

Entrez Gene:

9759 HDAC4 Related reagents

Synonyms	KIAA0288
Specificity	Mouse anti Human histone deacetylase 4 antibody recognizes the histone deacetylase 4 (HDAC4), also known as histone deacetylase A.
	Histones play a critical role in transcriptional regulation, cell cycle progression, and developmental events. Histone acetylation/deacetylation alters chromosome structure and affects transcription factor access to DNA. Encoded by the HDAC4 gene, histone deacetylase 4 belongs to class II of the histone deacetylase/acuc/apha family. It possesses histone deacetylase activity and represses transcription when tethered to a promoter. This protein does not bind DNA directly, but through transcription factors MEF2C and MEF2D. It seems to interact in a multiprotein complex with RbAp48 and HDAC3 (provided by RefSeq, Jul 2008).
	Mouse anti Human histone deacetylase 4 antibody detects a band of 140 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti histone deacetylase 4 detects a band of approximately 140 kDa in Jurkat cell lysates
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400μg Jurkat lysate lyophilized in RIPA buffer.
Lysate Reconstitution	 If using DDT reconstitute the lyophilized lysate with 190μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 10μl 2M DTT. If using BME reconstitute the lyophilized lysate with 180μl DI H₂O, add 200μl 2x Laemmli Sample Buffer and 20μl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25μl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

 North & South
 Tel: +1 800 265 7376
 Worldwide
 Tel: +44 (0)1865 852 700
 Europe
 Tel: +49 (0) 89 8090 95 21

 America
 Fax: +1 919 878 3751
 Fax: +44 (0)1865 852 739
 Fax: +49 (0) 89 8090 95 50

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351158:190315'

Printed on 09 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint