

Datasheet: VMA00293KT

Description:	TISSUE FACTOR PATHWAY INHIBITOR ANTIBODY WITH CONTROL LYSATE
Specificity:	TISSUE FACTOR PATHWAY INHIBITOR
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	OTI4H9
Isotype:	lgG2a
Quantity:	2 Westerns

Product Details

Applications	This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> .				
		Yes	No	Not Determined	Suggested Dilution
	Western Blotting	-			1/1000
	PrecisionAb antibodies <u>application.</u> The antibod product has not been tes exclude its use in such pr sample type.	ly has bee ted for us	en validato e in a par	ed at the suggested di ticular technique this c	lution. Where this does not necessarily
Target Species	Human				
Product Form	Purified IgG - liquid				
Preparation	20µl Mouse monoclonal a supernatant	antibody p	ourified by	/ affinity chromatograp	hy from tissue culture
Buffer Solution	Phosphate buffered salin	е			
Preservative	0.09% Sodium Azide (Na	aN ₃)			
Stabilisers	1% Bovine Serum Album	0,			
	50% Glycerol				
Immunogen	Recombinant protein com pathway inhibitor (NP_00	•	•		nan tissue factor

External Database	
Links	UniProt:
	P10646 Related reagents
	Entrez Gene:
	7035 TFPI Related reagents
Synonyms	LACI, TFPI1
Specificity	Mouse anti Human tissue factor pathway inhibitor antibody recognizes tissue factor pathway inhibitor, also known as anti-convertin and extrinsic pathway inhibitor.
	Encoded by the TFPI gene, tissue factor pathway inhibitor is a protease inhibitor that regulates the tissue factor (TF)-dependent pathway of blood coagulation. The coagulation process initiates with the formation of a factor VIIa-TF complex, which proteolytically activates additional proteases (factors IX and X) and ultimately leads to the formation of a fibrin clot. The product of TFPI inhibits the activated factor X and VIIa-TF proteases in an autoregulatory loop. The encoded protein is glycosylated and predominantly found in the vascular endothelium and plasma in both free forms and complexed with plasma lipoproteins. Several alternatively spliced transcript variants of TFPI have been described, but the full-length nature of some of these variants has not been confirmed (provided by RefSeq, Jul 2008).
	The antibody has been extensively validated for western blotting using whole cell lysates.
Western Blotting	Anti tissue factor pathway inhibitor detects a band of approximately 40-45 kDa in HepG2 cell lysates
Instructions For Use	Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page.
Lysate Composition	400µg HepG2 lysate lyophilized in RIPA buffer.
Lysate Reconstitution	 If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.
Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.
	Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.

Acknowledgements	PrecisionAb [™] is a trademark of Bio-Rad Laboratories.
------------------	--

Health And Safety	Material Safety Datasheet documentation #10048 #10561 available at:
Information	Antibody (10048): https://www.bio-rad-antibodies.com/uploads/MSDS/10048.pdf
	Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u>

Regulatory For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG2a NEGATIVE CONTROL (MCA929)

North & South	Tel: +1 800 265 7376	Worldwide
America	Fax: +1 919 878 3751	
	Email: antibody_sales_us@bio-rad.	com

 Tel: +44 (0)1865 852 700
 Europe

 Fax: +44 (0)1865 852 739
 Email: antibody_sales_uk@bio-rad.com

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com

From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351688:190318'

Printed on 10 Feb 2021

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint