

Datasheet: VMA00310

Description:	MOUSE ANTI TRANSCRIPTIONAL ACTIVATOR Myb
Specificity:	TRANSCRIPTIONAL ACTIVATOR Myb
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Clone:	1279CT309.289.119
Isotype:	lgG1
Quantity:	100 μΙ

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Immunoprecipitation	-			
Western Blotting				1/1000

The PrecisionAb label is reserved for antibodies that meet the defined performance criteria within Bio-Rad's ongoing antibody validation programme. Click here to learn how we validate our PrecisionAb range. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependent on sample type.

Target Species	Human
Product Form	Purified IgG - liquid
Preparation	Mouse monoclonal antibody prepared by affinity chromatography on Protein G from ascites.
Buffer Solution	Phosphate buffered saline.
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃).
Approx. Protein Concentrations	IgG concentration 0.5 mg/ml.
Immunogen	KLH-conjugated synthetic peptide from the central region of human transcriptional activator Myb

External	Database
Links	

UniProt:

P10242 Related reagents

Entrez Gene:

4602 MYB Related reagents

Specificity

Mouse anti Human transcriptional activator Myb antibody recognizes transcriptional activator Myb, also known as c-myb protein (140 AA), c-myb10A CDS, c-myb13A CDS, c-myb14A CDS, c-myb8B CDS and proto-oncogene c-Myb.

Encoded by the MYB gene, transcriptional activator Myb is a member of the MYB family of transcription factor genes. The protein contains three domains, an N-terminal DNA-binding domain, a central transcriptional activation domain and a C-terminal domain involved in transcriptional repression. This protein plays an essential role in the regulation of hematopoiesis and may play a role in tumorigenesis. Alternative splicing results in multiple transcript variants (provided by RefSeq, Jun 2009).

Mouse anti Human transcriptional activator Myb antibody detects a band of 80 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting

Anti transcriptional activator Myb detects a band of approximately 80 kDa in Jurkat cell

lysates.

Storage

This product is shipped at ambient temperature.

Store undiluted at -20°C, avoiding repeated freeze thaw cycles.

Guarantee

12 months from date of despatch.

Acknowledgements

PrecisionAb is a trademark of Bio-Rad Laboratories.

Health And Safety Information

Material Safety Datasheet documentation #10040 available at:

https://www.bio-rad-antibodies.com/SDS/VMA00310

Antibody (10040)

Regulatory

For research purposes only.

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG1 NEGATIVE CONTROL (MCA928)

North & South Tel: +1 800 265 7376

America

Worldwide Fax: +1 919 878 3751

Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Europe

Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50

Email: antibody_sales_us@bio-rad.com

Email: antibody_sales_uk@bio-rad.com

Email: antibody_sales_de@bio-rad.com

Printed on 23 May 2025

© 2025 Bio-Rad Laboratories Inc | Legal | Imprint