

Datasheet: VMA00329KT

Description:	PSMC5 ANTIBODY WITH CONTROL LYSATE
Specificity:	PSMC5
Format:	Purified
Product Type:	PrecisionAb Monoclonal
Isotype:	IgG1
Quantity:	2 Westerns

Product Details

Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit www.bio-rad-antibodies.com/protocols.

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

PrecisionAb antibodies have been extensively [validated for the western blot application](#). The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

Target Species	Human
Species Cross Reactivity	Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information.
Product Form	Purified IgG - liquid
Preparation	20µl Mouse monoclonal antibody prepared by affinity chromatography on Protein G from tissue culture supernatant
Buffer Solution	Phosphate buffered saline
Preservative Stabilisers	0.09% Sodium Azide (NaN ₃)

Immunogen Recombinant protein corresponding to amino acids 1-130 of human PSMC5 expressed in *E. coli*

External Database Links

UniProt:

[P62195](#) [Related reagents](#)

Entrez Gene:

[5705](#) PSMC5 [Related reagents](#)

Synonyms SUG1

Specificity

Mouse anti Human PSMC5 antibody recognizes the 26S protease regulatory subunit 8, also known as 26S protease regulatory subunit 8, 26S proteasome AAA-ATPase subunit RPT6, MSUG1 protein, Tat-binding protein homolog 10, proteasome 26S ATPase subunit 5, proteasome 26S subunit ATPase 5, proteasome subunit p45, thyroid hormone receptor-interacting protein 1, thyroid receptor interactor 1.

The 26S proteasome is a multicatalytic proteinase complex with a highly ordered structure composed of 2 complexes, a 20S core and a 19S regulator. The 20S core is composed of 4 rings of 28 non-identical subunits; 2 rings are composed of 7 alpha subunits and 2 rings are composed of 7 beta subunits. The 19S regulator is composed of a base, which contains 6 ATPase subunits and 2 non-ATPase subunits, and a lid, which contains up to 10 non-ATPase subunits. Proteasomes are distributed throughout eukaryotic cells at a high concentration and cleave peptides in an ATP/ubiquitin-dependent process in a non-lysosomal pathway. An essential function of a modified proteasome, the immunoproteasome, is the processing of class I MHC peptides. PSMC5 gene encodes one of the ATPase subunits, a member of the triple-A family of ATPases which have a chaperone-like activity. In addition to participation in proteasome functions, this subunit may participate in transcriptional regulation since it has been shown to interact with the thyroid hormone receptor and retinoid X receptor-alpha. Two transcript variants encoding different isoforms have been found for PSMC5 (provided by RefSeq, Nov 2010).

Mouse anti Human PSMC5 detects a band of 46 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.

Western Blotting Anti PSMC5 detects a band of approximately 46 kDa in HEK293 cell lysates

Instructions For Use Please refer to the [PrecisionAb western blotting protocol](#). For additional information on secondary antibody dilution and exposure time see product web page.

Lysate Composition 400µg HEK293 lysate lyophilized in RIPA buffer.

Lysate Reconstitution - If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.
- If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.
Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb

formats please refer to the PrecisionAb western blotting protocol.

Storage	Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.
Guarantee	As supplied, 12 months from date of despatch.
Acknowledgements	PrecisionAb™ is a trademark of Bio-Rad Laboratories.
Health And Safety Information	Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf Lysate Material (10561): https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf
Regulatory	For research purposes only

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

Recommended Negative Controls

[MOUSE IgG1 NEGATIVE CONTROL \(MCA928\)](#)

North & South America	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio-rad.com	Worldwide	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio-rad.com	Europe	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M351394:190318'

Printed on 09 Feb 2021