

Datasheet: VMA00340KT

| Description: | SUMO-2 ANTIBODY WITH CONTROL LYSATE |
|---------------|-------------------------------------|
| Specificity: | SUMO-2 |
| Format: | Purified |
| Product Type: | PrecisionAb Monoclonal |
| Isotype: | lgG2b |
| Quantity: | 2 Westerns |

Product Details

| Applications | This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit <u>www.bio-rad-antibodies.com/protocols</u> . | | | | | |
|-----------------------------|--|-------------------|-------------|------------------------|-----------------------|--|
| | | Yes | No | Not Determined | Suggested Dilution | |
| | Western Blotting | • | | | 1/1000 | |
| | PrecisionAb antibodies have been extensively validated for the western blot | | | | | |
| | application. The antibody has been validated at the suggested dilution. Where this | | | | | |
| | product has not been tested for use in a particular technique this does not necessarily | | | | • | |
| | exclude its use in such p sample type. | rocedures | . Further c | optimization may be re | equired dependant on | |
| Target Species | Human | | | | | |
| Species Cross Reactivity | Reacts with: Mouse, Rat N.B. Antibody reactivity and working conditions may vary between species. Cross reactivity is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. | | | | | |
| Product Form | Purified IgG - liquid | | | | | |
| Preparation | 20µl Mouse monoclonal tissue culture supernatar | | repared b | y affinity chromatogra | phy on Protein G from | |
| Buffer Solution | Phosphate buffered salin | ie | | | | |
| Preservative Stabilisers | 0.09% Sodium Azide (Na | aN ₃) | | | | |

| Immunogen | Recombinant protein corresponding to amino acids 1-95 of human SUMO-2 expressed in <i>E. coli</i> | | | |
|----------------------------|---|--|--|--|
| External Database Links | UniProt: <u>P61956</u> <u>Related reagents</u> Entrez Gene: <u>6613</u> SUMO2 <u>Related reagents</u> | | | |
| Synonyms | SMT3B, SMT3H2 | | | |
| Specificity | Mouse anti Human SUMO-2 antibody recognizes the small ubiquitin-related modifier 2 (SUMO-2), also known as SMT3 homolog 2, SMT3 suppressor of mif two 3 homolog 2, sentrin 2, small ubiquitin-related modifier 2, ubiquitin-like protein SMT3A and ubiquitin-like protein SMT3B. | | | |
| | Encoded by the SUMO2 gene, SUMO-2 is a member of the SUMO (small ubiquitin-like modifier) protein family. It functions in a manner similar to ubiquitin in that it is bound to target proteins as part of a post-translational modification system. However, unlike ubiquitin which targets proteins for degradation, this protein is involved in a variety of cellular processes, such as nuclear transport, transcriptional regulation, apoptosis, and protein stability. It is not active until the last two amino acids of the carboxy-terminus have been cleaved off. Numerous pseudogenes have been reported for SUMO2. Alternate transcriptional splice variants, encoding different isoforms, have been characterized (provided by RefSeq, Jul 2008). | | | |
| | Mouse anti Human SUMO-2 antibody detects a band of 18 kDa. The antibody has been extensively validated for western blotting using whole cell lysates. | | | |
| Western Blotting | Anti SUMO-2 detects a band of approximately 18 kDa in HEK293 cell lysates | | | |
| Instructions For Use | Please refer to the <u>PrecisionAb western blotting protocol.</u> For additional information on secondary antibody dilution and exposure time see product web page. | | | |
| Lysate Composition | 400µg HEK293 lysate lyophilized in RIPA buffer. | | | |
| Lysate Reconstitution | If using DDT reconstitute the lyophilized lysate with 190µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT. If using BME reconstitute the lyophilized lysate with 180µl DI H₂O, add 200µl 2x Laemmli Sample Buffer and 20µl BME. Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol. | | | |
| Storage | Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles. | | | |
| | Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage. | | | |

| Guarantee | As supplied, 12 months from date of despatch. |
|----------------------------------|--|
| Acknowledgements | PrecisionAb™ is a trademark of Bio-Rad Laboratories. |
| Health And Safety Information | Material Safety Datasheet documentation #10040 #10561 available at: Antibody (10040): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf</u> Lysate Material (10561): <u>https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf</u> |
| Regulatory | For research purposes only |

Related Products

Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) HRP

Recommended Negative Controls

MOUSE IgG2b NEGATIVE CONTROL (MCA691)

| North & South America | Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: antibody_sales_us@bio | Worldwide | Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: antibody_sales_uk@bio | Europe -rad.com | Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: antibody_sales_de@bio-rad.com | | |
|---|--|-----------|--|--------------------|--|--|--|
| From March 15, 2021, we will no longer supply printed datasheets with our products. Look out for updates on how to access your digital version at bio-rad-antibodies.com 'M351623:190318' | | | | | | | |
| Printed on 09 Feb 2021 | | | | | | | |

© 2021 Bio-Rad Laboratories Inc | Legal | Imprint