

## Datasheet: VMA00373KT

<b>Description:</b>	CD167 ANTIBODY WITH CONTROL LYSATE
<b>Specificity:</b>	CD167
<b>Format:</b>	Purified
<b>Product Type:</b>	PrecisionAb Monoclonal
<b>Isotype:</b>	IgG2b
<b>Quantity:</b>	2 Westerns

### Product Details

#### Applications

This product has been reported to work in the following applications. This information is derived from testing within our laboratories, peer-reviewed publications or personal communications from the originators. Please refer to references indicated for further information. For general protocol recommendations, please visit [www.bio-rad-antibodies.com/protocols](http://www.bio-rad-antibodies.com/protocols).

	Yes	No	Not Determined	Suggested Dilution
Western Blotting	▪			1/1000

**PrecisionAb antibodies have been extensively [validated for the western blot application](#).** The antibody has been validated at the suggested dilution. Where this product has not been tested for use in a particular technique this does not necessarily exclude its use in such procedures. Further optimization may be required dependant on sample type.

<b>Target Species</b>	Human
<b>Product Form</b>	Purified IgG - liquid
<b>Preparation</b>	20µl Mouse monoclonal antibody purified by affinity chromatography on Protein G from tissue culture supernatant
<b>Buffer Solution</b>	Phosphate buffered saline
<b>Preservative Stabilisers</b>	0.09% Sodium Azide (NaN <sub>3</sub> )
<b>Immunogen</b>	Recombinant CD167
<b>External Database Links</b>	<b>UniProt:</b> <a href="#">Q08345</a> <a href="#">Related reagents</a>

**Entrez Gene:**

[780](#) [DDR1](#) [Related reagents](#)

---

<b>Synonyms</b>	CAK, EDDR1, NEP, NTRK4, PTK3A, RTK6, TRKE
-----------------	---

---

<b>Specificity</b>	<p><b>Mouse anti Human CD167 antibody</b> recognizes CD167, also known as CD167 antigen-like family member A, PTK3A protein tyrosine kinase 3A, cell adhesion kinase, epithelial discoidin domain-containing receptor 1 or DDR1, mammary carcinoma kinase 10, protein-tyrosine kinase RTK-6, tyrosine kinase DDR and tyrosine-protein kinase CAK.</p> <p>Receptor tyrosine kinases play a key role in the communication of cells with their microenvironment. These kinases mostly regulate cell growth, differentiation and metabolism. CD167 belongs to a subfamily of tyrosine kinase receptors activated by various types of collagen with homology to Dictyostelium discoideum protein discoidin I in their extracellular domain. Expression of this protein is restricted to epithelial cells, particularly in the kidney, lung, gastrointestinal tract, and brain. In addition, it has been shown to be significantly overexpressed in several human tumors. Alternatively spliced transcript variants encoding different isoforms have been described for CD167 (provided by RefSeq, Feb 2011).</p> <p>Mouse anti Human CD167 antibody detects a band of 125 kDa. The antibody has been extensively validated for western blotting using whole cell lysates.</p>
--------------------	--

---

<b>Western Blotting</b>	Anti CD167 detects a band of approximately 125 kDa in MCF7 cell lysates
-------------------------	---

---

<b>Instructions For Use</b>	Please refer to the <a href="#">PrecisionAb western blotting protocol</a> . For additional information on secondary antibody dilution and exposure time see product web page.
-----------------------------	---

---

<b>Lysate Composition</b>	400µg MCF7 lysate lyophilized in RIPA buffer
---------------------------	--

---

<b>Lysate Reconstitution</b>	<p>- If using DDT reconstitute the lyophilized lysate with 190µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 10µl 2M DTT.</p> <p>- If using BME reconstitute the lyophilized lysate with 180µl DI H<sub>2</sub>O, add 200µl 2x Laemmli Sample Buffer and 20µl BME.</p> <p>Heat at 95°C for 5 minutes. For 10 well mini gels load 25µl. For other gel and comb formats please refer to the PrecisionAb western blotting protocol.</p>
------------------------------	--

---

<b>Storage</b>	<p>Antibody: Store undiluted at -20°C, avoiding repeated freeze thaw cycles.</p> <p>Lysate: Store lyophilized lysate at -20°C. After reconstitution aliquot and store at -20°C for up to 3 months or at -80°C for longer term storage.</p>
----------------	--

---

<b>Guarantee</b>	As supplied, 12 months from date of despatch
------------------	--

---

<b>Acknowledgements</b>	PrecisionAb™ is a trademark of Bio-Rad Laboratories
-------------------------	---

---

<b>Health And Safety</b>	Material Safety Datasheet documentation #10040 #10561 available at:
--------------------------	---

**Information** Antibody (10040): <https://www.bio-rad-antibodies.com/uploads/MSDS/10040.pdf>  
Lysate Material (10561): <https://www.bio-rad-antibodies.com/uploads/MSDS/10561.pdf>

---

**Regulatory** For research purposes only

---

## Related Products

### Recommended Secondary Antibodies

Goat Anti Mouse IgG (H/L) (STAR207...) [HRP](#)

### Recommended Negative Controls

[MOUSE IgG2b NEGATIVE CONTROL \(MCA691\)](#)

<b>North &amp; South America</b>	Tel: +1 800 265 7376 Fax: +1 919 878 3751 Email: <a href="mailto:antibody_sales_us@bio-rad.com">antibody_sales_us@bio-rad.com</a>	<b>Worldwide</b>	Tel: +44 (0)1865 852 700 Fax: +44 (0)1865 852 739 Email: <a href="mailto:antibody_sales_uk@bio-rad.com">antibody_sales_uk@bio-rad.com</a>	<b>Europe</b>	Tel: +49 (0) 89 8090 95 21 Fax: +49 (0) 89 8090 95 50 Email: <a href="mailto:antibody_sales_de@bio-rad.com">antibody_sales_de@bio-rad.com</a>
----------------------------------	---	------------------	---	---------------	---

From March 15, 2021, we will no longer supply printed datasheets with our products.  
Look out for updates on how to access your digital version at [bio-rad-antibodies.com](https://www.bio-rad-antibodies.com)

'M350952:190315'

**Printed on 09 Feb 2021**

---

© 2021 Bio-Rad Laboratories Inc | [Legal](#) | [Imprint](#)